

A Study of $B^0 \rightarrow \rho^+ \rho^-$ Decays and Constraints on the CKM Angle α .

B. Aubert, M. Bona, D. Boutigny, Y. Karyotakis, J. P. Lees, V. Poireau, X. Prudent, V. Tisserand, and A. Zghiche
Laboratoire de Physique des Particules, IN2P3/CNRS et Université de Savoie, F-74941 Annecy-Le-Vieux, France

J. Garra Tico and E. Grauges
Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

L. Lopez and A. Palano
Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

G. Eigen, I. Ofte, B. Stugu, and L. Sun
University of Bergen, Institute of Physics, N-5007 Bergen, Norway

G. S. Abrams, M. Battaglia, D. N. Brown, J. Button-Shafer, R. N. Cahn, Y. Groyzman, R. G. Jacobsen,
 J. A. Kadyk, L. T. Kerth, Yu. G. Kolomensky, G. Kukartsev, D. Lopes Pegna, G. Lynch, L. M. Mir,
 T. J. Orimoto, M. Pripstein, N. A. Roe, M. T. Ronan,* K. Tackmann, and W. A. Wenzel
Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

P. del Amo Sanchez, C. M. Hawkes, and A. T. Watson
University of Birmingham, Birmingham, B15 2TT, United Kingdom

T. Held, H. Koch, B. Lewandowski, M. Pelizaeus, T. Schroeder, and M. Steinke
Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

W. N. Cottingham and D. Walker
University of Bristol, Bristol BS8 1TL, United Kingdom

D. J. Asgeirsson, T. Cuhadar-Donszelmann, B. G. Fulsom,
 C. Hearty, N. S. Knecht, T. S. Mattison, and J. A. McKenna
University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

A. Khan, M. Saleem, and L. Teodorescu
Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

V. E. Blinov, A. D. Bukin, V. P. Druzhinin, V. B. Golubev, A. P. Onuchin,
 S. I. Serednyakov, Yu. I. Skovpen, E. P. Solodov, and K. Yu Todyshev
Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

M. Bondioli, S. Curry, I. Eschrich, D. Kirkby, A. J. Lankford, P. Lund, M. Mandelkern, E. C. Martin, and D. P. Stoker
University of California at Irvine, Irvine, California 92697, USA

S. Abachi and C. Buchanan
University of California at Los Angeles, Los Angeles, California 90024, USA

S. D. Foulkes, J. W. Gary, F. Liu, O. Long, B. C. Shen, and L. Zhang
University of California at Riverside, Riverside, California 92521, USA

H. P. Paar, S. Rahatlou, and V. Sharma
University of California at San Diego, La Jolla, California 92093, USA

J. W. Berryhill, C. Campagnari, A. Cunha, B. Dahmes, T. M. Hong, D. Kovalskyi, and J. D. Richman
University of California at Santa Barbara, Santa Barbara, California 93106, USA

T. W. Beck, A. M. Eisner, C. J. Flacco, C. A. Heusch, J. Kroseberg, W. S. Lockman,
T. Schalk, B. A. Schumm, A. Seiden, D. C. Williams, M. G. Wilson, and L. O. Winstrom
University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

E. Chen, C. H. Cheng, A. Dvoretzskii, F. Fang, D. G. Hitlin, I. Narsky, T. Piatenko, and F. C. Porter
California Institute of Technology, Pasadena, California 91125, USA

G. Mancinelli, B. T. Meadows, K. Mishra, and M. D. Sokoloff
University of Cincinnati, Cincinnati, Ohio 45221, USA

F. Blanc, P. C. Bloom, S. Chen, W. T. Ford, J. F. Hirschauer, A. Kreisel, M. Nagel,
U. Nauenberg, A. Olivas, J. G. Smith, K. A. Ulmer, S. R. Wagner, and J. Zhang
University of Colorado, Boulder, Colorado 80309, USA

A. M. Gabareen, A. Soffer, W. H. Toki, R. J. Wilson, F. Winklmeier, and Q. Zeng
Colorado State University, Fort Collins, Colorado 80523, USA

D. D. Altenburg, E. Feltresi, A. Hauke, H. Jasper, J. Merkel, A. Petzold, B. Spaan, and K. Wacker
Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany

T. Brandt, V. Klose, H. M. Lacker, W. F. Mader, R. Nogowski,
J. Schubert, K. R. Schubert, R. Schwierz, J. E. Sundermann, and A. Volk
Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

D. Bernard, G. R. Bonneaud, E. Latour, V. Lombardo, Ch. Thiebaux, and M. Verderi
Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, F-91128 Palaiseau, France

P. J. Clark, W. Gradl, F. Muheim, S. Playfer, A. I. Robertson, and Y. Xie
University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

M. Andreotti, D. Bettoni, C. Bozzi, R. Calabrese, A. Cecchi, G. Cibinetto, P. Franchini,
E. Luppi, M. Negrini, A. Petrella, L. Piemontese, E. Prencipe, and V. Santoro
Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy

F. Anulli, R. Baldini-Ferrolì, A. Calcaterra, R. de Sangro, G. Finocchiaro,
S. Pacetti, P. Patteri, I. M. Peruzzi,[†] M. Piccolo, M. Rama, and A. Zallo
Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy

A. Buzzo, R. Contri, M. Lo Vetere, M. M. Macri, M. R. Monge,
S. Passaggio, C. Patrignani, E. Robutti, A. Santroni, and S. Tosi
Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy

K. S. Chaisanguanthum, M. Morii, and J. Wu
Harvard University, Cambridge, Massachusetts 02138, USA

R. S. Dubitzky, J. Marks, S. Schenk, and U. Uwer
Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany

D. J. Bard, P. D. Dauncey, R. L. Flack, J. A. Nash, M. B. Nikolich, and W. Panduro Vazquez
Imperial College London, London, SW7 2AZ, United Kingdom

P. K. Behera, X. Chai, M. J. Charles, U. Mallik, N. T. Meyer, and V. Ziegler
University of Iowa, Iowa City, Iowa 52242, USA

J. Cochran, H. B. Crawley, L. Dong, V. Eyges, W. T. Meyer, S. Prell, E. I. Rosenberg, and A. E. Rubin
Iowa State University, Ames, Iowa 50011-3160, USA

A. V. Gritsan, Z. J. Guo, and C. K. Lae

Johns Hopkins University, Baltimore, Maryland 21218, USA

A. G. Denig, M. Fritsch, and G. Schott

Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany

N. Arnaud, J. Béquilleux, M. Davier, G. Grosdidier, A. Höcker, V. Lepeltier, F. Le Diberder, A. M. Lutz, S. Pruvot,
S. Rodier, P. Roudeau, M. H. Schune, J. Serrano, V. Sordini, A. Stocchi, W. F. Wang, and G. Wormser
*Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11,
Centre Scientifique d'Orsay, B. P. 34, F-91898 ORSAY Cedex, France*

D. J. Lange and D. M. Wright

Lawrence Livermore National Laboratory, Livermore, California 94550, USA

C. A. Chavez, I. J. Forster, J. R. Fry, E. Gabathuler, R. Gamet,
D. E. Hutchcroft, D. J. Payne, K. C. Schofield, and C. Touramanis
University of Liverpool, Liverpool L69 7ZE, United Kingdom

A. J. Bevan, K. A. George, F. Di Lodovico, W. Menges, and R. Sacco
Queen Mary, University of London, E1 4NS, United Kingdom

G. Cowan, H. U. Flaecher, D. A. Hopkins, P. S. Jackson, T. R. McMahon, F. Salvatore, and A. C. Wren
University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom

D. N. Brown and C. L. Davis

University of Louisville, Louisville, Kentucky 40292, USA

J. Allison, N. R. Barlow, R. J. Barlow, Y. M. Chia, C. L. Edgar, G. D. Lafferty, T. J. West, and J. I. Yi
University of Manchester, Manchester M13 9PL, United Kingdom

J. Anderson, C. Chen, A. Jawahery, D. A. Roberts, G. Simi, and J. M. Tuggle
University of Maryland, College Park, Maryland 20742, USA

G. Blaylock, C. Dallapiccola, S. S. Hertzbach, X. Li, T. B. Moore, E. Salvati, and S. Saremi
University of Massachusetts, Amherst, Massachusetts 01003, USA

R. Cowan, P. H. Fisher, G. Sciolla, S. J. Sekula, M. Spitznagel, F. Taylor, and R. K. Yamamoto
Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA

S. E. Mclachlin, P. M. Patel, and S. H. Robertson
McGill University, Montréal, Québec, Canada H3A 2T8

A. Lazzaro and F. Palombo

Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy

J. M. Bauer, L. Cremaldi, V. Eschenburg, R. Godang, R. Kroeger, D. A. Sanders, D. J. Summers, and H. W. Zhao
University of Mississippi, University, Mississippi 38677, USA

S. Brunet, D. Côté, M. Simard, P. Taras, and F. B. Viaud
Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7

H. Nicholson

Mount Holyoke College, South Hadley, Massachusetts 01075, USA

G. De Nardo, F. Fabozzi,[‡] L. Lista, D. Monorchio, and C. Sciacca
Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy

M. A. Baak, G. Raven, and H. L. Snoek

NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands

C. P. Jessop and J. M. LoSecco
University of Notre Dame, Notre Dame, Indiana 46556, USA

G. Benelli, L. A. Corwin, K. K. Gan, K. Honscheid, D. Hufnagel, H. Kagan, R. Kass,
 J. P. Morris, A. M. Rahimi, J. J. Regensburger, R. Ter-Antonyan, and Q. K. Wong
Ohio State University, Columbus, Ohio 43210, USA

N. L. Blount, J. Brau, R. Frey, O. Igonkina, J. A. Kolb, M. Lu,
 R. Rahmat, N. B. Sinev, D. Strom, J. Strube, and E. Torrence
University of Oregon, Eugene, Oregon 97403, USA

N. Gagliardi, A. Gaz, M. Margoni, M. Morandin, A. Pompili,
 M. Posocco, M. Rotondo, F. Simonetto, R. Stroili, and C. Voci
Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy

E. Ben-Haim, H. Briand, J. Chauveau, P. David, L. Del Buono, Ch. de la Vaissière,
 O. Hamon, B. L. Hartfiel, Ph. Leruste, J. Malclès, J. Ocariz, and A. Perez
*Laboratoire de Physique Nucléaire et de Hautes Energies,
 IN2P3/CNRS, Université Pierre et Marie Curie-Paris6,
 Université Denis Diderot-Paris7, F-75252 Paris, France*

L. Gladney
University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA

M. Biasini, R. Covarelli, and E. Manoni
Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy

C. Angelini, G. Batignani, S. Bettarini, G. Calderini, M. Carpinelli, R. Cenci, A. Cervelli, F. Forti, M. A. Giorgi,
 A. Lusiani, G. Marchiori, M. A. Mazur, M. Morganti, N. Neri, E. Paoloni, G. Rizzo, and J. J. Walsh
Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy

M. Haire
Prairie View A&M University, Prairie View, Texas 77446, USA

J. Biesiada, P. Elmer, Y. P. Lau, C. Lu, J. Olsen, A. J. S. Smith, and A. V. Telnov
Princeton University, Princeton, New Jersey 08544, USA

E. Baracchini, F. Bellini, G. Cavoto, A. D'Orazio, D. del Re, E. Di Marco, R. Faccini, F. Ferrarotto, F. Ferroni,
 M. Gaspero, P. D. Jackson, L. Li Gioi, M. A. Mazzoni, S. Morganti, G. Piredda, F. Polci, F. Renga, and C. Voena
Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy

M. Ebert, H. Schröder, and R. Waldi
Universität Rostock, D-18051 Rostock, Germany

T. Adye, G. Castelli, B. Franek, E. O. Olaiya, S. Ricciardi, W. Roethel, and F. F. Wilson
Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom

R. Aleksan, S. Emery, M. Escalier, A. Gaidot, S. F. Ganzhur, P. F. Giraud,
 G. Hamel de Monchenault, W. Kozanecki, M. Legendre, G. Vasseur, Ch. Yèche, and M. Zito
DSM/Daphnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France

X. R. Chen, H. Liu, W. Park, M. V. Purohit, and J. R. Wilson
University of South Carolina, Columbia, South Carolina 29208, USA

M. T. Allen, D. Aston, R. Bartoldus, P. Bechtle, N. Berger, R. Claus, J. P. Coleman, M. R. Convery, J. C. Dingfelder,
 J. Dorfan, G. P. Dubois-Felsmann, D. Dujmic, W. Dunwoodie, R. C. Field, T. Glanzman, S. J. Gowdy,
 M. T. Graham, P. Grenier, C. Hast, T. Hryn'ova, W. R. Innes, M. H. Kelsey, H. Kim, P. Kim, D. W. G. S. Leith,

S. Li, S. Luitz, V. Luth, H. L. Lynch, D. B. MacFarlane, H. Marsiske, R. Messner, D. R. Muller, C. P. O'Grady, A. Perazzo, M. Perl, T. Pulliam, B. N. Ratcliff, A. Roodman, A. A. Salnikov, R. H. Schindler, J. Schwiening, A. Snyder, J. Stelzer, D. Su, M. K. Sullivan, K. Suzuki, S. K. Swain, J. M. Thompson, J. Va'vra, N. van Bakel, A. P. Wagner, M. Weaver, W. J. Wisniewski, M. Wittgen, D. H. Wright, A. K. Yarritu, K. Yi, and C. C. Young
Stanford Linear Accelerator Center, Stanford, California 94309, USA

P. R. Burchat, A. J. Edwards, S. A. Majewski, B. A. Petersen, and L. Wilden
Stanford University, Stanford, California 94305-4060, USA

S. Ahmed, M. S. Alam, R. Bula, J. A. Ernst, V. Jain, B. Pan, M. A. Saeed, F. R. Wappler, and S. B. Zain
State University of New York, Albany, New York 12222, USA

W. Bugg, M. Krishnamurthy, and S. M. Spanier
University of Tennessee, Knoxville, Tennessee 37996, USA

R. Eckmann, J. L. Ritchie, A. M. Ruland, C. J. Schilling, and R. F. Schwitters
University of Texas at Austin, Austin, Texas 78712, USA

J. M. Izen, X. C. Lou, and S. Ye
University of Texas at Dallas, Richardson, Texas 75083, USA

F. Bianchi, F. Gallo, D. Gamba, and M. Pelliccioni
Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy

M. Bomben, L. Bosisio, C. Cartaro, F. Cossutti, G. Della Ricca, L. Lanceri, and L. Vitale
Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy

V. Azzolini, N. Lopez-March, F. Martinez-Vidal, D. A. Milanes, and A. Oyanguren
IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain

J. Albert, Sw. Banerjee, B. Bhuyan, K. Hamano, R. Kowalewski, I. M. Nugent, J. M. Roney, and R. J. Sobie
University of Victoria, Victoria, British Columbia, Canada V8W 3P6

J. J. Back, P. F. Harrison, T. E. Latham, G. B. Mohanty, and M. Pappagallo[§]
Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom

H. R. Band, X. Chen, S. Dasu, K. T. Flood, J. J. Hollar,
P. E. Kutter, Y. Pan, M. Pierini, R. Prepost, S. L. Wu, and Z. Yu
University of Wisconsin, Madison, Wisconsin 53706, USA

H. Neal
Yale University, New Haven, Connecticut 06511, USA

We present results from an analysis of $B^0 \rightarrow \rho^+ \rho^-$ decays using $(383.6 \pm 4.2) \times 10^6 B\bar{B}$ pairs collected by the *BABAR* detector at the PEP-II asymmetric-energy *B* Factory at SLAC. The measurements of the $B^0 \rightarrow \rho^+ \rho^-$ branching fraction, longitudinal polarization fraction f_L , and the *CP*-violating parameters S_{long} and C_{long} are:

$$\begin{aligned} \mathcal{B}(B^0 \rightarrow \rho^+ \rho^-) &= (25.5 \pm 2.1(\text{stat})_{-3.9}^{+3.6}(\text{syst})) \times 10^{-6}, \\ f_L &= 0.992 \pm 0.024(\text{stat})_{-0.013}^{+0.026}(\text{syst}), \\ S_{\text{long}} &= -0.17 \pm 0.20(\text{stat})_{-0.06}^{+0.05}(\text{syst}), \\ C_{\text{long}} &= 0.01 \pm 0.15(\text{stat}) \pm 0.06(\text{syst}). \end{aligned}$$

We determine the unitarity triangle angle α , using an isospin analysis of $B \rightarrow \rho\rho$ decays. One of the two solutions, $\alpha = [73.1, 117.0]^\circ$ at 68% CL is compatible with standard model-based fits of existing data. Constraints on the unitarity triangle are also evaluated using an *SU*(3) symmetry based approach.

I. INTRODUCTION

Charge conjugation-parity (CP) violation was first seen in the decay of neutral kaons [1]. It was shown some forty years ago that CP violation is a necessary but insufficient condition required to generate a net baryon anti-baryon asymmetry in the universe [2]. The standard model (SM) of electroweak interactions describes CP violation as a consequence of a complex phase in the three-generation Cabibbo-Kobayashi-Maskawa (CKM) quark-mixing matrix [3, 4]:

$$V_{\text{CKM}} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}. \quad (1)$$

V_{CKM} describes the couplings of the u , c and t quarks to d , s and b quarks, which are mediated by the exchange of a W boson. In B -meson decays the CP violating parameters of the SM are most directly related to the angles and sides of the so-called Unitarity Triangle (UT), shown in Fig. 1. The angles α , β , and γ are defined as

$$\alpha \equiv \arg[-V_{td}V_{tb}^*/V_{ud}V_{ub}^*], \quad (2)$$

$$\beta \equiv \arg[-V_{cd}V_{cb}^*/V_{td}V_{tb}^*], \quad (3)$$

$$\gamma \equiv \arg[-V_{ud}V_{ub}^*/V_{cd}V_{cb}^*]. \quad (4)$$

Any non-trivial phase in V_{ij} is CP violating. CP violating phases originating from the CKM matrix are related to weak interactions, and therefore referred to as weak phases. In the Wolfenstein convention [5], the angle γ is the phase of V_{ub} , β is the phase of V_{td} , and α is the phase difference between V_{ub} and V_{td} constrained to satisfy $\alpha = \pi - \beta - \gamma$ through the unitarity of V_{CKM} .




FIG. 1: The Unitarity Triangle in the $\bar{\rho}$ - $\bar{\eta}$ plane.

In Fig. 1 the UT is shown in the complex $(\bar{\rho}, \bar{\eta})$ plane, where the apex is given by the phase-convention independent definition [6]:

$$\bar{\rho} + i\bar{\eta} \equiv -\frac{V_{ud}V_{ub}^*}{V_{cd}V_{cb}^*}. \quad (5)$$

The quest to understand CP violation remains, despite its observation in the B meson system by both the BABAR [7] and Belle experiments [8], since the SM does not, through the CKM phase, incorporate enough CP violation to explain the current matter-antimatter asymmetry [9, 10]. The CKM angle β is measured to a 1° precision with $b \rightarrow c\bar{c}s$ decays [11, 12] and is consistent with current predictions [13–15]. A significant deviation from results of SM-based fits of existing data for any of the UT angles would be a clear indication of new physics [16, 17].

In principle, direct experimental measurements of α can be made from decays that proceed mainly through a $\bar{b} \rightarrow u\bar{u}d$ tree diagram such as $B^0 \rightarrow \rho^+\rho^-$, $\rho^\pm\pi^\mp$, $\pi^+\pi^-$ and $a_1\pi$ [18, 19]. Interference between the direct decay and decay after $B^0\bar{B}^0$ mixing in each of these decays results in a time-dependent decay-rate asymmetry that is sensitive to the angle α . Figure 2 shows the leading order tree and gluonic penguin loop contributions to the decay $B^0 \rightarrow \rho^+\rho^-$. The presence of penguin contributions complicates the extraction of α from these decays. Using isospin relations [20], measurements of the $B^+ \rightarrow \rho^+\rho^0$ [21, 22] and $B^0 \rightarrow \rho^0\rho^0$ [23] branching fractions show that the penguin contribution in $B \rightarrow \rho\rho$ is smaller than the leading tree diagram. The use of $SU(3)$ flavor symmetry to increase the precision on the weak phase constrained using $B^0 \rightarrow \rho^+\rho^-$ and $B^+ \rightarrow K^{*0}\rho^+$ decays has also been proposed [24]. Both of these approaches are discussed in Section X. Section X and Ref. [25] describe a number of possible sources of theoretical uncertainty.




FIG. 2: Tree (top) and gluonic penguin (bottom) diagrams contributing to the process $B^0 \rightarrow \rho^+\rho^-$. The penguin contribution coming from the diagram with a top quark in the loop dominates as contributions from processes with u and c quarks are suppressed.

In $B^0 \rightarrow \rho^+\rho^-$ decays, a spin 0 particle (the B^0 meson) decays into two spin 1 particles ($\rho^\pm$ mesons), as shown in Fig. 3. Subsequently each $\rho^\pm$ meson decays into a $\pi^\pm\pi^0$ pair. As a result, the CP analysis of $B^0 \rightarrow \rho^+\rho^-$ is complicated by the presence of one amplitude with longitudinal polarization and two amplitudes with transverse polarization. The longitudinal amplitude is CP -even, while

*Deceased

[†]Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

[‡]Also with Università della Basilicata, Potenza, Italy

[§]Also with IPPP, Physics Department, Durham University, Durham DH1 3LE, United Kingdom

the transverse amplitudes contain CP -even and CP -odd states. The decay is observed to be dominated by the longitudinal polarization [26, 27], with a fraction f_L defined as the fraction of the helicity zero state in the decay. Integrating over the angle between the ρ decay planes ϕ , the angular decay rate is

$$\frac{d^2\Gamma}{\Gamma d\cos\theta_1 d\cos\theta_2} = \frac{9}{4} \left[f_L \cos^2\theta_1 \cos^2\theta_2 + \frac{1}{4}(1 - f_L) \sin^2\theta_1 \sin^2\theta_2 \right], \quad (6)$$

where the helicity angles $\theta_{i=1,2}$ are the angles between the π^0 momentum and the direction opposite to that of the B^0 in the ρ rest frame.




FIG. 3: A schematic of the decay of a B meson via two ρ mesons to a four pion final state. The ρ meson final states are shown in their rest frames, and ϕ is the angle between the decay planes of the ρ mesons.

We identify (tag) the initial flavor of the reconstructed B candidate, B_{rec} , using information from the other B meson, B_{tag} , in the event. The decay rate $f_+(f_-)$ for a neutral B meson decaying into a CP eigenstate, where the B_{tag} is a B^0 ($\bar{B}^0$), is given by:

$$f_{\pm}(\Delta t) = \frac{e^{-|\Delta t|/\tau_{B^0}}}{4\tau_{B^0}} \left\{ 1 \pm \eta_f [S \sin(\Delta m_d \Delta t) - C \cos(\Delta m_d \Delta t)] \right\}. \quad (7)$$

Here Δt is the proper time difference between the decay of the B_{rec} and B_{tag} mesons in an event, τ_{B^0} is the mean B^0 lifetime, Δm_d is the B^0 - $\bar{B}^0$ mixing frequency [28], and η_f is the CP eigenvalue of the decay. This assumes that there is no difference between the B^0 and $\bar{B}^0$ lifetimes, $\Delta\Gamma = 0$. The parameters S and C describe B -mixing induced and direct CP violation, respectively. For the longitudinal polarization $\eta_f = +1$, $S = S_{\text{long}}$ and $C =$

C_{long} , are defined as

$$S_{\text{long}} = \frac{2\text{Im}\lambda_{CP}}{1 + |\lambda_{CP}|^2}, \quad (8)$$

$$C_{\text{long}} = \frac{1 - |\lambda_{CP}|^2}{1 + |\lambda_{CP}|^2}, \quad (9)$$

where $\lambda_{CP} = \frac{q}{p} \frac{\bar{A}}{A}$ [29], q and p are complex constants that relate the B meson flavor eigenstates to the mass eigenstates, and $\bar{A}/A$ is the ratio of amplitudes of the decay of a $\bar{B}^0$ or B^0 to the final state under study. CP violation is probed by studying the time-dependent decay-rate asymmetry

$$\mathcal{A}(\Delta t) = \frac{\Gamma(\Delta t) - \bar{\Gamma}(\Delta t)}{\Gamma(\Delta t) + \bar{\Gamma}(\Delta t)}, \quad (10)$$

where $\Gamma(\bar{\Gamma})$ is the decay-rate for B^0 ($\bar{B}^0$) tagged events. This asymmetry has the form

$$\mathcal{A}(\Delta t) = S \sin(\Delta m_d \Delta t) - C \cos(\Delta m_d \Delta t). \quad (11)$$

The transverse polarization is an admixture of CP -even ($\eta_f = +1$) and CP -odd ($\eta_f = -1$) parts, where each CP eigenstate has a decay-rate distribution of the form of Eq. (7). As the longitudinal polarization dominates this decay, we describe the CP admixture of the transverse polarization with common parameters S_{tran} and C_{tran} . We set $S_{\text{tran}} = C_{\text{tran}} = 0$, and vary these parameters when evaluating systematic uncertainties.

In the absence of penguin contributions in $B^0 \rightarrow \rho^+ \rho^-$, $S_{\text{long}} = \sin 2\alpha$ and $C_{\text{long}} = 0$. The presence of penguin contributions with different weak phases to the tree level amplitude shifts the experimentally measurable parameter α_{eff} away from the value of α . In the presence of penguin contributions $\alpha_{\text{eff}} = \alpha + \delta\alpha$, where

$$S_{\text{long}} = \sqrt{1 - C_{\text{long}}^2} \sin 2\alpha_{\text{eff}}, \quad (12)$$

and C_{long} can be non-zero.

For $B^+ \rightarrow \rho^+ \rho^0$ one measures a time-integrated charge asymmetry, which is analogous to Eq. (10) and is defined as

$$A_{CP} = \frac{\bar{N} - N}{\bar{N} + N}, \quad (13)$$

where N ($\bar{N}$) is the number of B ($\bar{B}$) decays to this final state. Table I summarizes the most recent measurements of the complete set of experimental inputs needed to extract α from the $B \rightarrow \rho\rho$ decays.

In this article, we present an update of the time-dependent analysis of $B^0 \rightarrow \rho^+ \rho^-$ and measurement of the CKM angle α reported in Ref. [26] and branching fraction reported in Ref. [30]. Improvements to the analysis reported here include an increase in data sample analyzed, a tighter constraint on the proper time difference between the two B meson decays in selected events, an improved algorithm to determine the flavor of B mesons, a modified multivariate analyzer for background suppression, and an improved description of the signal and background model.

TABLE I: Measurements of the $B \rightarrow \rho\rho$ decays. Branching fractions ($\mathcal{B}$) are in units of 10^{-6} . The column marked $C_{\text{long}}(A_{CP})$ denotes the time dependent charge asymmetry C_{long} in the case of the $B^0 \rightarrow \rho^+\rho^-$ decay, and the time integrated charge asymmetry A_{CP} in the case of $B^\pm \rightarrow \rho^\pm\rho^0$.

Mode	Expt. (luminosity fb^{-1})	$\mathcal{B}$ ($\times 10^{-6}$)	f_L	$C_{\text{long}}(A_{CP})$	S_{long}
$B^0 \rightarrow \rho^+\rho^-$	BABAR [30] (82)	$30 \pm 4 \pm 5$	—	—	—
$B^0 \rightarrow \rho^+\rho^-$	BABAR [26] (210)	—	$0.978 \pm 0.014^{+0.021}_{-0.029}$	$-0.03 \pm 0.18 \pm 0.09$	$-0.33 \pm 0.24^{+0.08}_{-0.14}$
$B^0 \rightarrow \rho^+\rho^-$	Belle [27] (253)	$22.8 \pm 3.8^{+2.3}_{-2.6}$	$0.941^{+0.034}_{-0.040} \pm 0.030$	—	—
$B^0 \rightarrow \rho^+\rho^-$	Belle [31] (535)	—	—	$-0.16 \pm 0.21 \pm 0.08^a$	$0.19 \pm 0.30 \pm 0.08$
$B^\pm \rightarrow \rho^\pm\rho^0$	BABAR [21] (210)	$16.8 \pm 2.2 \pm 2.3$	$0.905 \pm 0.042^{+0.023}_{-0.027}$	$-0.12 \pm 0.13 \pm 0.10$	—
$B^\pm \rightarrow \rho^\pm\rho^0$	Belle [22] (78)	$31.7 \pm 7.1^{+3.8}_{-6.7}$	$0.95 \pm 0.11 \pm 0.02$	$0.00 \pm 0.22 \pm 0.03$	—
$B^0 \rightarrow \rho^0\rho^0$	BABAR [23] (349)	$1.07 \pm 0.33 \pm 0.19$	$0.87 \pm 0.13 \pm 0.04$	—	—

^aBelle Collaboration uses the opposite sign convention for C_{long} .

II. THE DATASET AND BABAR DETECTOR

The results presented in this paper are based on data collected with the *BABAR* detector at the PEP-II asymmetric e^+e^- storage ring [32] operating at the Stanford Linear Accelerator Center. At PEP-II, 9.0 GeV electrons and 3.1 GeV positrons are collided at a center-of-mass energy of 10.58 GeV which corresponds to the mass of the $\Upsilon(4S)$ resonance. The asymmetric energies result in a boost from the laboratory to the center-of-mass (CM) frame of $\beta\gamma \approx 0.56$. The dataset analyzed has an integrated luminosity of 349 fb^{-1} corresponding to $(383.6 \pm 4.2) \times 10^6 B\bar{B}$ pairs recorded at the $\Upsilon(4S)$ resonance (on-peak). An additional 27.2 fb^{-1} of data were recorded about 40 MeV below this energy (off-peak) for the study of continuum background, where light quarks are produced in the process $e^+e^- \rightarrow q\bar{q}$ ($q = u, d, s, c$).

The *BABAR* detector is described in detail elsewhere [33]. Surrounding the interaction point is a five double-sided layer silicon vertex tracker (SVT) which measures the impact parameters of charged particle tracks in both the plane transverse to, and along the beam direction. A 40-layer drift chamber (DCH) surrounds the SVT and provides measurements of the momenta for charged particles. Both the SVT and DCH are surrounded by a solenoid magnet, that provides a 1.5 T magnetic field. Charged hadron identification is achieved through measurements of particle energy-loss in the tracking system and the Cherenkov angle obtained from a detector of internally reflected Cherenkov light. A CsI(Tl) electromagnetic calorimeter (EMC) provides photon detection, electron identification, and π^0 reconstruction. Finally, the instrumented flux return of the magnet allows discrimination of muons from pions. For the most recent 134.0 fb^{-1} of data, a portion of the resistive plate chambers constituting the muon system has been replaced by limited streamer tubes [34–36].

We use a right-handed coordinate system with the z axis along the electron beam direction and the y axis upward, with the origin at the nominal beam interaction

point. Unless otherwise stated, kinematic quantities are calculated in the laboratory rest frame. The other reference frame which we commonly use is the CM frame of the colliding electrons and positrons.

We use Monte Carlo (MC) simulated events generated using the GEANT4 [37] based *BABAR* simulation.

III. RECONSTRUCTION OF B CANDIDATES

A. Photon and π^0 reconstruction

Photons are reconstructed from localized energy deposits in the EMC that are not associated with a charged track. We require photon candidates to have an energy greater than 50 MeV, and a lateral shower profile [38] to be consistent with the photon hypothesis. We reconstruct neutral pions from combinations of two distinct photon candidates where the invariant $\gamma\gamma$ mass $m_{\gamma\gamma}$ satisfies $0.10 < m_{\gamma\gamma} < 0.16 \text{ GeV}/c^2$.

B. $\rho^\pm$ reconstruction

We combine reconstructed π^0 mesons with charged tracks that are consistent with the $\pi^\pm$ hypothesis to form $\rho^\pm$ candidates. The invariant mass $m_{\pi^\pm\pi^0}$ of the reconstructed $\rho^\pm$ is required to lie between 0.5 and $1.0 \text{ GeV}/c^2$, to select events in the vicinity of the ρ resonance. We require that the helicity angle of each ρ meson satisfies $-0.90 < \cos\theta_i < 0.98$. This selection criteria suppresses continuum and B backgrounds.

C. B^0 reconstruction and event selection

We combine two oppositely charged ρ candidates to form the B candidate B_{rec} . All particles in the rest of the event (ROE) are combined to form the other B meson in the event B_{tag} . In addition to the two charged tracks

in the B_{rec} , we require that there is at least one track in the B_{tag} .

In order to suppress potential backgrounds from $e^+e^- \rightarrow e^+e^-, \mu^+\mu^-$ events, we require the second-to-zeroth Fox-Wolfgram moment R_2 [39] of the event to be less than 0.98. Continuum events are the dominant background which is reduced by requiring the absolute value of the cosine of the angle between the B_{rec} thrust axis (TB) and that of the ROE (TR) to satisfy $|\cos(TB, TR)| < 0.8$. We retain 17.1% and 20.1% of longitudinal and transverse signal, respectively, by requiring the aforementioned selection criteria.

We calculate $\Delta t = \Delta z / \beta \gamma c$ from the measured separation Δz between the B_{rec} and B_{tag} vertices [40]. We determine the B_{rec} vertex from the two charged-pion tracks in its decay. The B_{tag} decay vertex is obtained by fitting the other tracks in the event, with constraints from the B_{rec} momentum and the beam-spot location. The RMS resolution on Δt is 1.1 ps. We only use events that satisfy $|\Delta t| < 15$ ps and for which the error on Δt ($\sigma_{\Delta t}$) is less than 2.5 ps.

Signal events are identified using two kinematic variables, the difference ΔE between the CM energy of the B_{rec} , E_B^* , and $\sqrt{s}/2$,

$$\Delta E = E_B^* - \sqrt{s}/2, \quad (14)$$

and the beam-energy-substituted mass,

$$m_{ES} = \sqrt{(s/2 + \mathbf{p}_i \cdot \mathbf{p}_B)^2 / E_i^2 - \mathbf{p}_B^2}, \quad (15)$$

where $\sqrt{s}$ is the total CM energy. The B_{rec} momentum $\mathbf{p}_B$ and four-momentum of the initial state ($E_i, \mathbf{p}_i$) are defined in the laboratory frame. We accept candidates that satisfy $5.25 < m_{ES} < 5.29$ GeV/ c^2 and $-0.12 < \Delta E < 0.15$ GeV. An asymmetric ΔE selection is used in order to reduce backgrounds from higher-multiplicity B decays. The resolution on m_{ES} is dominated by the knowledge of the energy of the e^+ and e^- beams, and the resolution on ΔE is dominated by the reconstruction performance of the EMC.

After the selection criteria mentioned above have been applied, the average number of candidates per event is approximately 1.5. In events with more than one candidate, we select the candidate that minimizes the sum of $(m_{\gamma\gamma} - m_{\pi^0})^2$ where m_{π^0} is the true π^0 mass [28]. In 0.3% of events, the same π^0 mesons are used by multiple B candidates. In such an event we randomly select the candidate to keep.

IV. CONTINUUM BACKGROUND SUPPRESSION

In addition to the cuts on $\cos\theta_i$, R_2 , and $|\cos(TB, TR)|$ that directly remove background events, we use an artificial neural network in order to discriminate between continuum background and signal events.

For this purpose we combine the following eight variables into a single output, $\mathcal{N}$.

- The coefficients, L_0, L_2 , split into sums over the ROE for neutral and charged particles; $L_{0,n}, L_{2,n}$ and $L_{0,c}, L_{2,c}$. The coefficients are defined as $L_k = \sum p_j |\cos(\psi_j)|^k$, where $k = 0, 2$, p_j is the particle momentum and ψ_j is the angle of the particle direction relative to the thrust axis of the B candidate. Both p_j and ψ_j are defined in the CM frame.
- $|\cos(B, Z)|$, the absolute value of the cosine of the angle between the direction of the B and z axis in the CM frame. This variable follows a sine squared distribution for $B\bar{B}$ events, whereas it is almost uniform for $q\bar{q}$.
- $|\cos(TB, TR)|$. This variable, previously defined in Section III C, is strongly peaked at unity for $q\bar{q}$ events. $B\bar{B}$ events are more isotropic as the B mesons are produced close to the kinematic threshold.
- $|\cos(TB, Z)|$, the absolute value of the cosine of the angle between the B thrust and the z axis.
- The scalar sum of the transverse momenta p_t in the ROE. This sum includes neutral and charged particles.

The distributions of these input variables are shown in Fig. 4. Figure 5 shows $\mathcal{N}$ for signal MC simulated events and continuum background samples (off-peak data) and the efficiency for signal and $q\bar{q}$ background as a function of cut on the minimum value of $\mathcal{N}$. We require $\mathcal{N}$ to be greater than -0.4 . Note that later, we use this variable in the maximum likelihood fit described in Section VII.

The samples used to train the neural net were correctly reconstructed MC simulated events and off-peak data. To avoid over-training, we used an independent sample of these data (*i.e.* distinct from the sample used for the training) to check the performance of the network. The training is stopped when the separation between the signal and continuum background is stable.

V. MIS-RECONSTRUCTED SIGNAL AND SELECTION EFFICIENCY

Mis-reconstructed signal candidates, referred to as self-cross-feed (SCF) signal, may pass the selection requirements even if one or more of the pions assigned to the $\rho^+\rho^-$ state belongs to the other B in the event. These SCF candidates constitute 50.7% (27.9%) of the accepted longitudinally (transversely) polarized signal. The majority of SCF events have both charged pions from the $\rho^+\rho^-$ final state, and unbiased CP information. These correct (right) track SCF events are denoted by RT SCF. There is a SCF component (13.8% of the signal) where at least one track in B_{rec} is from the ROE. These wrong




FIG. 4: The input variables used in training the neural network. The solid line represents the signal training sample (MC simulated events) and the dashed line represents the continuum background (off-peak data). The distributions shown are (in order left to right; top to bottom) $L_{0,n}$, $L_{0,c}$, $L_{2,n}$, $L_{2,c}$, $\cos(B, Z)$, $\cos(B, TR)$, $\cos(TB, Z)$, and the sum of the p_t in the ROE.




FIG. 5: The left plot shows the distribution of $\mathcal{N}$ for off-peak data (dashed) and longitudinally polarized signal MC simulated events (solid). The right plot shows the signal efficiency (solid) and continuum background efficiency (dashed) distribution as a function of cut on the minimum value of $\mathcal{N}$.

track (WT) events are used to determine the signal yield N_{signal} and f_L but have biased CP information, and are treated separately for the CP result. A systematic error is assigned to the CP results from this type of signal event. The total selection efficiency for longitudinally (transversely) polarized signal is 7.7% (10.5%).

VI. B MESON FLAVOR TAGGING

A key ingredient in the measurement of time-dependent CP asymmetries is to determine whether at the time of decay of the B_{tag} , corresponding to $\Delta t = 0$, the B_{rec} was a B^0 or a $\bar{B}^0$. This ‘flavor tagging’ is achieved with the analysis of the decay products of the recoiling B meson B_{tag} . The overwhelming majority of

B mesons decay to a final state that is flavor-specific, i.e. only accessible from either a B^0 or a $\bar{B}^0$, but not from both. The purpose of the flavor tagging algorithm is to determine the flavor of B_{tag} with the highest possible efficiency ϵ_{tag} and lowest possible probability w of assigning a wrong flavor to B_{tag} . It is not necessary to fully reconstruct B_{tag} in order to determine its flavor. In the presence of a finite mistag probability w , the CP asymmetry is reduced by a dilution factor $1 - 2w$, so Eq. (7) becomes

$$f_{\pm}(\Delta t) = \frac{e^{-|\Delta t|/\tau_{B^0}}}{4\tau_{B^0}} \left\{ 1 \pm (1 - 2w) \times \eta_f [S \sin(\Delta m_d \Delta t) - C \cos(\Delta m_d \Delta t)] \right\}. \quad (16)$$

The figure of merit for the performance of the tagging algorithm is the effective tagging efficiency

$$Q = \epsilon_{\text{tag}}(1 - 2w)^2, \quad (17)$$

which is related to the statistical uncertainty σ in the coefficients S and C through

$$\sigma \propto \frac{1}{\sqrt{Q}}. \quad (18)$$

We use a neural network based technique [11, 40] that isolates primary leptons, kaons and pions from B decays to final states containing D^* mesons, and high momentum charged particles from B decays, to determine the flavor of the B_{tag} . The output of this algorithm is divided into seven mutually-exclusive categories. These are (in order of decreasing signal purity) Lepton, Kaon I, Kaon II, Kaon-Pion, Pion, Other and Untagged. The performance of this algorithm is determined using fully reconstructed neutral B decays to $D^{(*)-}(\pi^+, \rho^+, a_1^+)$ and is summarized in Table II. The categories assigned correspond to events with leptons, kaons and pions in the decay products of B_{tag} . The Untagged category of events contain no flavor information and therefore carry no weight in the time-dependent analysis. In addition, these events are not considered useful for the branching fraction measurement since they are dominated by continuum background. The total Q of this algorithm is $32.7 \pm 0.7\%$.

VII. LIKELIHOOD FIT METHOD

On applying the selection criteria described above, we obtain a sample of 37424 events that enter the fit. These events are dominated by backgrounds from $q\bar{q}$ (81.4%) and $B\bar{B}$ (16.6%) events. The remaining 2% of events are considered as signal. We distinguish between the following components in the fit:

(i) correctly reconstructed signal,

TABLE II: Tagging efficiency ϵ , average mistag fraction w , mistag fraction difference between B^0 and $\bar{B}^0$ tagged events Δw , and Q for $B^0 \rightarrow \rho^+ \rho^-$ events.

Category	ϵ_{tag} (%)	w (%)	Δw (%)	Q (%)
Lepton	8.2 ± 0.1	3.2 ± 0.5	-0.2 ± 0.8	7.2 ± 0.2
Kaon I	11.3 ± 0.1	3.7 ± 0.7	1.1 ± 1.2	9.7 ± 0.3
Kaon II	17.3 ± 0.2	14.2 ± 0.7	-0.9 ± 1.1	8.8 ± 0.3
Kaon-Pion	13.4 ± 0.1	20.8 ± 0.8	0.5 ± 1.3	4.6 ± 0.3
Pion	13.8 ± 0.2	30.6 ± 0.8	4.1 ± 1.3	2.1 ± 0.2
Other	9.4 ± 0.1	40.1 ± 1.0	2.3 ± 1.5	0.4 ± 0.1
Untagged	26.8 ± 0.2	50.0 ± 0.0	—	0.0 ± 0.0
Total				32.7 ± 0.7

(ii) SCF signal, split into RT and WT parts,

(iii) charm $B^\pm$ backgrounds ($b \rightarrow c$),

(iv) charm B^0 backgrounds ($b \rightarrow c$),

(v) charmless B^0 backgrounds,

(vi) charmless $B^\pm$ backgrounds,

(vii) continuum background.

The dominant B backgrounds come from components (iii) and (iv). The signal, continuum and B background models are described in Sections VII A, VII B, and VII C, respectively.

We use an unbinned, extended maximum likelihood (ML) fit to extract N_{signal} , f_L , S_{long} , and C_{long} . The likelihood function incorporates the following previously defined eight discriminating variables to distinguish signal from the backgrounds: m_{ES} , ΔE , Δt , $\mathcal{N}$, and the m_k and $\cos \theta_k$ values of the two ρ mesons, where $k = 1, 2$. For each of the aforementioned components j we construct a probability density function (PDF) that is the product of one-dimensional PDFs for each of the variables. The PDFs do not account for all possible correlations among the discriminating variables and we account for possible biases as a result of neglecting these correlations as discussed in Section VIII. For each event i , the PDFs can be written as

$$\mathcal{P}_j^i = \mathcal{P}_j(m_{ES}^i) \mathcal{P}_j(\Delta E^i) \mathcal{P}_j(\Delta t^i) \mathcal{P}_j(\mathcal{N}^i) \times \mathcal{P}_j(m_1^i) \mathcal{P}_j(m_2^i) \mathcal{P}_j(\cos \theta_1^i) \mathcal{P}_j(\cos \theta_2^i). \quad (19)$$

We determine the PDFs for signal and each of the B background components from MC distributions for each discriminating variable. The PDFs for the continuum background are determined from on-peak and off-peak data. The likelihood function is

$$\mathcal{L} = \frac{\exp(-\sum_j n_j)}{N!} \prod_i \sum_j n_j \mathcal{P}_j^i, \quad (20)$$

where n_j are the event yields of hypotheses j (determined from the ML fit) and N is the observed number of events

in the sample. The normalized exponent takes into account Poisson fluctuations in the total number of events. We simultaneously fit events in the mutually-exclusive flavor tag categories listed in Table II, excluding events in the **Untagged** category. There are six continuum background yields, one for each flavor tag category, and we use a single yield for the signal and each of the B backgrounds, while accounting for the relative fractions of events expected in each of the flavor tag categories. The fit results are obtained by minimizing the value of $-\ln \mathcal{L}$.

A. Signal model

The signal has contributions from true and SCF longitudinally (long) and transversely (tran) polarized events. In addition to this, the longitudinally polarized SCF signal PDF is further sub-divided into the two categories of mis-reconstructed signal: RT and WT SCF signal, and all signal PDFs are sub-divided according to the B_{tag} flavor tag category. The total signal PDF is given by

$$\begin{aligned} \mathcal{P}_{\text{signal}} = & f_L \left([1 - f_{\text{RTSCF}}^{\text{long}} - f_{\text{WTSCF}}^{\text{long}}] \mathcal{P}_{\text{true}}^{\text{long}} \right. \\ & \left. + f_{\text{RTSCF}}^{\text{long}} \mathcal{P}_{\text{RTSCF}}^{\text{long}} + f_{\text{WTSCF}}^{\text{long}} \mathcal{P}_{\text{WTSCF}}^{\text{long}} \right) \\ & + (1 - f_L) ([1 - f_{\text{SCF}}^{\text{tran}}] \mathcal{P}_{\text{true}}^{\text{tran}} + f_{\text{SCF}}^{\text{tran}} \mathcal{P}_{\text{SCF}}^{\text{tran}}), \end{aligned} \quad (21)$$

where $f_{\text{RT(WT)SCF}}^{\text{long}}$ is the fraction of longitudinally polarized RT(WT) SCF signal and $f_{\text{SCF}}^{\text{tran}}$ is the fraction of transversely polarized SCF signal. The PDFs $\mathcal{P}_{\text{RT(WT)SCF}}^{\text{long}}$ and $\mathcal{P}_{\text{SCF}}^{\text{tran}}$ are defined accordingly. In order to extract the observable f_L from the fit, we account for the different reconstruction efficiencies for longitudinally and transversely polarized signal using

$$f_L = \frac{f_L^{\text{obs}}}{R + f_L^{\text{obs}}(1 - R)}, \quad (22)$$

where f_L^{obs} is the fraction of signal from longitudinally polarized events in the data, and R is the ratio of longitudinal to transverse polarized signal efficiencies.

All of the SCF PDFs are further sub-divided into parts that contain correctly reconstructed and mis-reconstructed ρ mesons. Four sub-divisions are used, corresponding to both mesons being correctly reconstructed (TT), one being correctly reconstructed and the other being mis-reconstructed (TF or FT), and both mesons being mis-reconstructed (FF). Here the first (second) ρ meson is positively (negatively) charged. The SCF PDFs

have the form

$$\begin{aligned} \mathcal{P} = & \mathcal{P}(\mathcal{N})\mathcal{P}(\Delta t)[\\ & f_{TT}\mathcal{P}_{TT}(m_{ES}, \Delta E, \cos \theta_1, m_1, \cos \theta_2, m_2) + \\ & f_{TF}\mathcal{P}_{TF}(m_{ES}, \Delta E, \cos \theta_1, m_1, \cos \theta_2, m_2) + \\ & f_{FT}\mathcal{P}_{FT}(m_{ES}, \Delta E, \cos \theta_1, m_1, \cos \theta_2, m_2) + \\ & f_{FF}\mathcal{P}_{FF}(m_{ES}, \Delta E, \cos \theta_1, m_1, \cos \theta_2, m_2)], \end{aligned} \quad (23)$$

where T and F are correctly and mis-reconstructed ρ mesons, respectively, f_{TT} , f_{TF} , f_{FT} , f_{FF} are the fractions of each type of mis-reconstructed event, and the PDFs for m_{ES} , ΔE , and the helicity and mass for each reconstructed ρ meson are the products of one dimensional PDFs. The signal PDFs for $\mathcal{N}$ and Δt have different shapes according to the B_{tag} flavor tag category assigned to an event.

The longitudinally polarized true and RT SCF signal m_{ES} distribution are described by a Gaussian with a power law tail [41–43] which takes the following form

$$\begin{aligned} f(m) \propto & \exp\left(-\frac{(m - m_0)^2}{2\sigma^2}\right), \quad m > m_0 - \xi\sigma, \\ & \propto \frac{n^n \exp(-\xi^2/2)}{\left(\frac{\xi(m_0 - m)}{\sigma} + n - \xi^2\right)^n}, \quad m \leq m_0 - \xi\sigma, \end{aligned} \quad (24)$$

where m_0 is the mean, σ is the width of the Gaussian part, and the parameters ξ and n describe the exponential tail. The remaining PDFs for the signal m_{ES} distributions are the sum of a Gaussian and a Gaussian with an exponential tail. The signal ΔE distributions for the longitudinally polarized SCF events are described by first and second order polynomials, with the exception of TT events, which are modeled with the sum of a second order polynomial and a Gaussian. The other ΔE PDFs are described by the sum of Gaussian and a Gaussian with an exponential tail. The $\mathcal{N}$ distributions are modeled using smoothed histograms of MC simulated events. The signal Δt distribution is described by Eq. (16) convolved with a triple Gaussian resolution function given by

$$\begin{aligned} \mathcal{R}_{\text{sig}}(\Delta t, \sigma_{\Delta t}) = & f_{\text{core}}G(\Delta t, \mu_{\text{core}}\sigma_{\Delta t}, \sigma_{\text{core}}\sigma_{\Delta t}) + \\ & f_{\text{tail}}G(\Delta t, \mu_{\text{tail}}\sigma_{\Delta t}, \sigma_{\text{tail}}\sigma_{\Delta t}) + \\ & f_{\text{outlier}}G(\Delta t, \mu_{\text{outlier}}, \sigma_{\text{outlier}}), \end{aligned} \quad (25)$$

where G is a Gaussian with means μ_l and width σ_l for $l = \text{core, tail, and outlier}$. The parameters σ_{tail} , σ_{outlier} and μ_{outlier} are set to 3.0 ps, 8.0 ps and 0.0 ps, respectively. The remaining parameters of the resolution function are determined from signal MC simulated events scaled by the differences between data and MC simulated events for large samples of exclusively reconstructed B decays to $D^{(*)-}(\pi^+, \rho^+, a_1^+)$ final states described in Section VI. The values of the mean and width of the core Gaussian contribution to the resolution function depend on the flavor tagging category of an event. There are

three signal Δt PDFs, one for the true and RT SCF longitudinally polarized signal, one for the WT SCF longitudinally polarized signal and one for the transversely polarized signal. Each of these PDFs has distinct values of S and C as described in Sections I and V. The WT SCF longitudinally polarized signal has a different resolution function with respect to the rest of the signal. The $\cos\theta_i$ distribution for true signal events is described by the expected cosine squared or sine squared distribution multiplied by a polynomial acceptance function. The $\cos\theta_i$ PDFs for T ρ mesons in longitudinally (transversely) polarized SCF signal are described by sixth order polynomials (smoothed histograms of MC simulated events). The $\cos\theta_i$ PDFs for mis-reconstructed ρ mesons are described by smoothed histograms of MC simulated events. The longitudinally polarized signal ρ mass PDFs are described using relativistic Breit-Wigners and third order polynomials for T and F ρ contributions, respectively. The transversely polarized signal SCF ρ mass PDF distributions are described using smoothed histograms of MC simulated events.

In addition to N_{signal} , f_L , S_{long} , and C_{long} , we determine the mean of the core Gaussian part of the m_{ES} and ΔE distributions from the fit. We obtain parameters consistent with expectations from MC simulated events.

B. Continuum background model

The continuum background PDFs are also sub-divided according to the mis-reconstruction of ρ mesons in the final state and have the form

$$\mathcal{P} = \mathcal{P}(m_{ES})\mathcal{P}(\Delta E)\mathcal{P}(\mathcal{N})\mathcal{P}(\Delta t) \left[\begin{aligned} &f_{TT}\mathcal{P}_{TT}(\cos\theta_1, m_1, \cos\theta_2, m_2) + \\ &f_{TF}\mathcal{P}_{TF}(\cos\theta_1, m_1, \cos\theta_2, m_2) + \\ &f_{FT}\mathcal{P}_{FT}(\cos\theta_1, m_1, \cos\theta_2, m_2) + \\ &f_{FF}\mathcal{P}_{FF}(\cos\theta_1, m_1, \cos\theta_2, m_2) \end{aligned} \right], \quad (26)$$

where helicity and mass PDFs for each reconstructed ρ meson are the products of one dimensional PDFs.

The continuum distribution for m_{ES} is described by a phase-space-motivated distribution [44] with the following form

$$f(x) \propto x\sqrt{1-x^2} \cdot \exp[\xi(1-x^2)] \cdot \theta(m_{ES}), \quad (27)$$

where $\theta(m_{ES}) = 1$ for $m_{ES} \leq \sqrt{s}/2$ and $\theta(m_{ES}) = 0$ for $m_{ES} > \sqrt{s}/2$ and $x = 2m_{ES}/\sqrt{s}$. The ΔE and $\mathcal{N}$ shapes are modeled with third and fourth order polynomials, respectively. The parameters of the m_{ES} , ΔE and $\mathcal{N}$ shapes are allowed to vary in the fit to the on-peak data. The continuum ρ mass distribution is described using a relativistic Breit-Wigner and a third order polynomial PDF for T and F ρ contributions, respectively. The

$\cos\theta_i$ distribution is described by a third order polynomial. The continuum Δt distribution has a prompt lifetime component convolved with a triple Gaussian resolution function. The parameters of the ρ mass and helicity distributions are obtained from a fit to the off-peak data, and the remaining parameters are determined in the fit.

C. B background model

1. Charm B backgrounds

Combinatorial events from $b \rightarrow c$ transitions are the dominant B backgrounds. These components have shapes similar to continuum and do not peak in the signal region for the discriminating variables. The functional form used for the PDFs of these background components is given by Eq. (26). We parameterize the m_{ES} and ΔE distributions of these backgrounds using the phase-space-motivated distribution of Eq. (27), and a third order polynomial, respectively. The ρ mass distribution is described using a relativistic Breit-Wigner and a third order polynomial PDF for T and F ρ contributions, respectively. The remaining PDFs are described using smoothed histograms of MC simulated events. Each of the PDF parameters for these backgrounds are determined from samples of MC simulated events, and the yields of these components are determined in the fit. When studying systematic uncertainties we modify the PDF used for Δt so that it has a form similar to the signal which uses the resolution function of Eq. (25). An effective lifetime that is smaller than τ_{B^0} is used to account for mis-reconstruction of these events and the finite charm meson lifetime. The value of this parameter is obtained by fitting MC simulated events for this category of events.

2. Charmless B backgrounds

Some of the charmless B backgrounds have PDFs similar to the signal for one or more of the discriminating variables, so it is important to correctly account for such events in the fit. We consider the 20 exclusive and 2 inclusive components of this type of B background listed in Table III. If a charmless B background decay contributes an event yield more than 1% of the expected signal yield, we model that mode exclusively. The remaining 140 charmless background components considered were combined with the appropriate weightings to form the neutral and charged inclusive charmless B background components. Where possible we use branching fractions from existing measurements. Where measurements are not available, we have either tried to use $SU(2)$ and $SU(3)$ flavor symmetries to relate an unmeasured decay to a measured one, or where this is not possible, we have assigned a branching fraction of 10×10^{-6} to a decay mode. An uncertainty of 100% is assumed on all

TABLE III: The components of charmless B backgrounds considered, along with the branching fraction ($\mathcal{B}$) and number of events (N_{bg}) expected in the selected data sample, where $\dagger$ indicates a longitudinally polarized final state, and NR denotes a non-resonant final state. Where appropriate, branching fractions of these decays are quoted, including the branching fraction of sub-decay modes and measured or expected values of f_L .

Decay Mode	$\mathcal{B} \times 10^{-6}$	N_{bg}
$B^+ \rightarrow K^+ \pi^+ \pi^-$ [45, 46]	55 ± 3	11 ± 1
$B^+ \rightarrow \pi^+ \pi^0 \pi^0$	10 ± 10	15 ± 15
$B^+ \rightarrow \pi^0 \rho^+$ [47–49]	11 ± 2	73 ± 10
$B^+ \rightarrow \rho^+ \rho^0$ [21, 22] $\dagger$	17 ± 3	71 ± 12
$B^+ \rightarrow \rho^+ \pi^+ \pi^-$	10 ± 10	9 ± 8
$B^+ \rightarrow K_2^*(1770) \rho$	10 ± 10	9 ± 9
$B^+ \rightarrow K_0^* \pi$ [45, 46, 49]	11 ± 1	9 ± 1
$B^+ \rightarrow a_1^+ \pi^0$	20 ± 20	60 ± 60
$B^+ \rightarrow \rho^+ \pi^0 \pi^0$	10 ± 10	12 ± 12
$B^+ \rightarrow a_1^0 (\rho^+ \pi^-) \rho^+ \dagger$	8 ± 8	13 ± 13
$B^+ \rightarrow a_1^0 \pi^+$	20 ± 20	49 ± 49
$B^+ \rightarrow \text{charmless}$	–	59 ± 15
$B^0 \rightarrow \pi^+ \pi^- \pi^0$ (NR)	30 ± 3	23 ± 23
$B^0 \rightarrow \rho^\pm \pi^\mp$ [49, 50]	24 ± 3	42 ± 4
$B^0 \rightarrow \pi^0 \pi^0 K_S^0$	23 ± 23	15 ± 15
$B^0 \rightarrow a_1^\pm (\rho^0 \pi^\pm) \pi^\mp$ [51]	20 ± 2	7 ± 1
$B^0 \rightarrow \rho^\mp \pi^\pm \pi^0$	10 ± 10	45 ± 45
$B^0 \rightarrow K_2^*(1770) \rho$	10 ± 10	8 ± 8
$B^0 \rightarrow a_1^\pm (\rho^\pm \pi^0) \rho^\mp$ [52] $\dagger$	16 ± 16	43 ± 43
$B^0 \rightarrow a_1^\pm (\rho^0 \pi^\pm) \rho^\mp$ [52] $\dagger$	16 ± 15	9 ± 8
$B^0 \rightarrow a_1^\pm (\rho^\pm \pi^0) \pi^\mp$ [51]	40 ± 4	102 ± 9
$B^0 \rightarrow \text{charmless}$	–	88 ± 22

extrapolated branching fractions. We assume that the a_1 meson decays into a three pion final state via $\rho\pi$. The decay $B^0 \rightarrow a_1^0 \pi^0$ is penguin dominated and is expected to have a small branching fraction, as is the case for the penguin dominated decays $B^0 \rightarrow \rho^0 \pi^0$ and $B^0 \rightarrow \rho^0 \rho^0$. As a result, we have neglected possible contributions from this potential source of background. Although we don't see evidence for $B^0 \rightarrow \rho^\mp \pi^\pm \pi^0$, we do consider this channel as a potential source of background. Contribution from $B^0 \rightarrow 4\pi$ is expected to be even smaller and is neglected. When considering the systematic uncertainty arising from interference between signal and other $\pi^+ \pi^- \pi^0 \pi^0$ final states, we assume that the non-resonant $\pi^+ \pi^- \pi^0 \pi^0$ final state has the same branching fraction as that of $B^0 \rightarrow \rho^\mp \pi^\pm \pi^0$, where $\mathcal{B}(B^0 \rightarrow \rho^\mp \pi^\pm \pi^0)$ is calculated from the yield determined in the $B^0 \rightarrow \rho^+ \rho^-$ nominal fit. This is discussed further in the Appendix.

The functional form used for the PDFs of these charmless B background components is given by Eq. (26). The ΔE distributions are described by third order polynomials, except for non-resonant $B^0 \rightarrow \rho^\mp \pi^\pm \pi^0$ which uses smoothed histograms of MC simulated events. The m_ρ distributions for true ρ mesons are parameterized using a relativistic Breit-Wigner, and the fake m_ρ (combinatorial $\pi^\pm \pi^0$) distribution is described using third order poly-

nomials. The remaining background shapes are described using smoothed histograms of MC simulated events. The yield for $B^0 \rightarrow \rho^\mp \pi^\pm \pi^0$ decays is allowed to vary in the fit. All other charmless background yields are fixed to expectations. This constraint is relaxed when studying possible sources of systematic uncertainties. When studying systematic uncertainties from possible CP violation in the B background, we modify the PDF used for Δt so that it has a form similar to the signal which uses the signal resolution function in Eq. (25).

VIII. LIKELIHOOD FIT VALIDATION

Before applying the fitting procedure to the data, we subject it to various tests. The aim of these tests are to verify that one can extract the signal observables N_{signal} , f_L , S_{long} , and C_{long} in a controlled way. Consistency of the likelihood fit is checked with ensembles of experiments simulated from the PDFs. The event yields are generated according to a Poisson distribution with mean n_j for each category in these ensembles. In each of these tests we verify that the values generated for the signal observables are reproduced with the expected resolution. The distribution of $-\ln \mathcal{L}$ for the ensemble of experiments in comparison to that obtained when fitting the data provides an additional, but limited, cross-check of the consistency.

The PDFs used in the likelihood do not account for all possible correlations among the discriminating variables. We account for possible biases as a result of neglecting these correlations by fitting ensembles of experiments obtained from samples of signal and the charmless B background MC simulated events combined with charm backgrounds and $q\bar{q}$ background events simulated from the PDFs. The MC simulated events used in these ensembles have these correlations modeled. We find a positive bias of 38.5 ± 5.3 events on N_{signal} , and a negative bias of 0.016 ± 0.002 on f_L , and we do not observe a significant bias on S_{long} and C_{long} .

As continuum events are the dominant background, we apply the fitting procedure to the off-peak data (after correcting for the difference in $\sqrt{s}$) to confirm that we do not find a fake signal in this control sample of events. We fit 790 off-peak events and extract signal and continuum yields of 8 ± 7 and 782 ± 28 events, respectively.

A blind analysis technique has been used for the extraction of S_{long} and C_{long} , where the actual values of these observables have been hidden by an offset. With the values of S_{long} and C_{long} hidden we perform the following fit cross-checks. We first verify that the uncertainties on the signal observables, and the value of $-\ln \mathcal{L}$ obtained from the fit to data, are compatible with the ensembles of simulated experiments described above. We then validate the stability of our results by introducing a variety of modifications to the fitting procedure. In addition to the nominal set of variables determined from data, we allow the physics parameters τ_{B^0} and Δm_d to

vary in turn. The signal observables do not change significantly when doing this, and the results obtained for τ_{B^0} and Δm_d are 1.72 ± 0.16 ps and $0.36 \pm 0.22 \hbar/\text{ps}$, respectively, consistent with the reported world average [28].

The **Lepton** and **Kaon l** tagged events have the highest signal purity, and lowest w probability, and dominate our CP results. We check that the results obtained from fitting only these categories of events are compatible with our nominal blind results. When doing this, we observe shifts of -0.08 (-0.05) on S_{long} (C_{long}) relative to the nominal result.

Given that there are a number of B backgrounds that contribute to the data that have yields fixed to expectations, we validate this assumption by allowing each fixed B background yield to vary in the fit. We obtain background yields consistent with our expectations, and observe the shifts on signal parameters listed in Table IV. A systematic uncertainty corresponding to the largest observed deviation is assigned on our results.

Once these checks have been completed, the fitting procedure is frozen and we extract the values of S_{long} and C_{long} by removing the offset.

TABLE IV: The shifts on signal yield ($\delta N(\text{signal})$), fraction of longitudinally polarized events (δf_L), S_{long} (δS_{long}), and C_{long} (δC_{long}) obtained when floating the yield of each fixed B background in turn. The symbol $\dagger$ indicates a longitudinally polarized final state, and NR denotes a non-resonant final state. The yields of decay modes not listed in this table are allowed to vary in the nominal fit.

Decay Mode	$\delta N(\text{signal})$	δf_L	δS_{long}	δC_{long}
$B^+ \rightarrow K^+ \pi^+ \pi^-$	3	-0.002	-0.002	-0.003
$B^+ \rightarrow \pi^+ \pi^0 \pi^0$	11	0.001	0.002	-0.001
$B^+ \rightarrow \pi^0 \rho^+$	9	0.000	0.001	-0.002
$B^+ \rightarrow \rho^+ \rho^0 \dagger$	-8	-0.001	-0.002	0.000
$B^+ \rightarrow \rho^+ \pi^+ \pi^-$	1	0.000	-0.001	-0.001
$B^+ \rightarrow K_2^*(1770) \rho$	-5	0.001	-0.001	-0.000
$B^+ \rightarrow K_0^* \pi$	-13	-0.001	-0.002	0.001
$B^+ \rightarrow a_1^+ \pi^0$	17	0.003	0.006	-0.001
$B^+ \rightarrow \rho^+ \pi^0 \pi^0$	-2	0.000	0.001	-0.002
$B^+ \rightarrow a_1^0(\rho^+ \pi^-) \rho^+ \dagger$	-34	-0.006	-0.028	-0.003
$B^+ \rightarrow a_1^0 \pi^+$	5	0.001	0.004	-0.001
$B^+ \rightarrow \text{charmless}$	-3	-0.001	-0.002	0.000
$B^0 \rightarrow \pi^+ \pi^- \pi^0$ (NR)	-4	0.000	0.000	0.000
$B^0 \rightarrow \rho^\pm \pi^\mp$	-5	0.000	-0.001	0.000
$B^0 \rightarrow \pi^0 \pi^0 K_S^0$	-25	-0.003	-0.009	-0.001
$B^0 \rightarrow a_1^\pm(\rho^0 \pi^\pm) \pi^\mp$	1	-0.002	-0.001	-0.002
$B^0 \rightarrow K_2^*(1770) \rho$	-31	-0.003	-0.009	-0.003
$B^0 \rightarrow a_1^\pm(\rho^\pm \pi^0) \rho^\mp \dagger$	-25	-0.002	-0.025	-0.004
$B^0 \rightarrow a_1^\pm(\rho^0 \pi^\pm) \rho^\mp \dagger$	11	0.000	0.006	0.000
$B^0 \rightarrow a_1^\pm(\rho^\pm \pi^0) \pi^\mp$	10	0.002	0.003	-0.002
$B^0 \rightarrow \text{charmless}$	1	0.000	0.000	-0.001

IX. RESULTS

A. Fit results

From the ML fit described above, we obtain the following results

$$\begin{aligned} N(\text{signal}) &= 729 \pm 60(\text{stat}), \\ f_L &= 0.992 \pm 0.024(\text{stat}), \\ S_{\text{long}} &= -0.17 \pm 0.20(\text{stat}), \\ C_{\text{long}} &= 0.01 \pm 0.15(\text{stat}), \end{aligned}$$

after correction for a $+38.5 \pm 5.3$ event fit bias (see Section VIII), a -76 event bias from SCF on the signal yield (see Section IX B) and a correction for a -0.016 ± 0.002 fit bias on f_L . The correlation between S_{long} and C_{long} is -0.035 . We discuss the origin of these fit biases in Section IX B. We calculate the branching fraction of this decay using $\mathcal{B} = N_{\text{signal}}/(\epsilon N_{B\bar{b}})$, where ϵ is the efficiency for signal corresponding to the observed f_L , and $N_{B\bar{b}}$ is the number of $B\bar{B}$ pairs analyzed. We obtain

$$\mathcal{B}(B^0 \rightarrow \rho^+ \rho^-) = (25.5 \pm 2.1(\text{stat})) \times 10^{-6}.$$

The $B^0 \rightarrow \rho^\pm \pi^\pm \pi^0$ background yield obtained from the fit is 9.2 ± 53.6 events. Figure 6 shows distributions of m_{ES} , ΔE , $\cos \theta_i$ and $m_{\pi^\pm \pi^0}$ for the **Lepton** and **Kaon l** tagged events with a loose requirement on $\mathcal{N}$. Relative to the total number of events in the fit, the plot of m_{ES} contains 15.6% of the signal and 1.1% of the total background. For the other plots there is an added constraint that $m_{ES} > 5.27 \text{ GeV}/c^2$; these requirements retain 13.9% of the signal and 0.4% of the total background. Figure 7 shows the Δt distribution for B^0 and $\bar{B}^0$ tagged events, as well as the time-dependent decay-rate asymmetry of Eq. (10). Here we apply the same selection criteria as in Fig. 6(b)-(d).


B. Systematic uncertainty studies

Table V lists the possible sources of systematic uncertainties on the values of the N_{signal} , f_L , S_{long} , and C_{long} that have been studied, and are described in the following.


- The uncertainty from PDF parameterisation is obtained by varying PDF shape parameters by $\pm 1\sigma$, in turn. The deviations obtained are added in quadrature to give the quoted uncertainty.
- The systematic uncertainty from the fraction of SCF events is obtained from the difference between the nominal fit result and that obtained when fitting the data and removing the SCF from the fit model. In the case of the signal yield we correct for half of the difference observed, and use 100% of the correction as an uncertainty. The uncertainty on the other signal observables comes from the difference observed between the two sets of fit results.

TABLE V: Summary of additive systematic uncertainty contributions on the signal yield N_{signal} , f_L , S_{long} and C_{long} .

Contribution	$\sigma(N_{\text{signal}})$	$\sigma(f_L)$	$\sigma(S_{\text{long}})$	$\sigma(C_{\text{long}})$
PDF parameterisation	+23 -41	+0.019 -0.005	+0.02 -0.04	0.03
SCF fraction	76	0.003	0.00	0.02
Control samples calibration	16	0.006	0.01	0.01
m_{ES} endpoint	12	0.001	0.00	0.01
B background normalization	+16 -20	+0.005 -0.002	0.01	0.00
floating B backgrounds	34	0.006	0.03	0.00
B background CP asymmetry	2	+0.001 -0.000	0.00	+0.02 -0.01
τ_B	+0 -1	+0.001 -0.000	0.00	0.00
Δm_d	+0 -1	+0.001 -0.000	0.00	0.00
tagging and dilution	+2 -16	+0.010 -0.001	0.00	0.01
transverse polarization CP asymmetry	+0 -9	+0.006 -0.000	0.01	0.01
Wrong track SCF CP asymmetry	+0 -3	+0.001 -0.000	0.01	0.01
DCSD decays	—	—	0.01	0.04
Interference	18	0.000	0.01	0.01
Fit Bias	19	0.008	0.02	0.02
SVT Alignment	—	—	0.01	0.01
Total	+94 -102	+0.03 -0.01	+0.05 -0.06	0.06

FIG. 6: The distributions for the highest purity tagged events for the variables (a) m_{ES} , (b) ΔE , (c) cosine of the ρ helicity angle and (d) $m_{\pi^\pm\pi^0}$. The dashed lines are the sum of backgrounds, and the solid lines are the full PDF.

- The kinematic endpoint position in m_{ES} is extracted from the fit. Changes in beam energy in the data can affect the endpoint position. To ac-

FIG. 7: The Δt distribution for a sample of events enriched in signal for (a) B^0 and (b) $\bar{B}^0$ tagged events. The dashed lines are the sum of backgrounds, and the solid lines are the sum of signal and backgrounds. The time-dependent CP asymmetry (see text) is shown in (c), where the curve is the measured asymmetry.

count for possible effects of this, we vary the kinematic endpoint position in m_{ES} by $\pm 0.45 \text{ MeV}/c^2$

which is determined from control samples of fully reconstructed B^0 decays.

- The uncertainty from the m_{ES} and ΔE widths is obtained from the observed shifts relative to our nominal result, when allowing these parameters to vary independently in the fit to data.
- We vary the B background normalization within expectations for each background in turn. The deviations obtained are added in quadrature to give the quoted uncertainty from this source.
- As the branching fractions of some of the B backgrounds are not well known, we assign an additional uncertainty coming from the maximum shifts obtained when allowing each of the fixed backgrounds to vary in turn in the fit to data.
- Additional uncertainties on the CP results come from possible CP violation in the B background. We replace the Δt PDFs used by each of the B backgrounds in turn by one resembling the signal. Charged B backgrounds can have non-zero values of A_{CP} , and neutral B backgrounds can have non-zero values of S and C . We use existing experimental constraints where possible, otherwise we allow for a CP asymmetry up to 10% in B decays to final states with charm, and up to 50% in B decays to charmless final states.
- The physics parameters $\tau_{B^0} = 1.530 \pm 0.009$ ps and $\Delta m_d = 0.507 \pm 0.005$ $\hbar$ /ps [28] are varied within the quoted uncertainty.
- The tagging and mistag fractions for signal and the B backgrounds are corrected for data/MC differences observed in samples of fully reconstructed hadronic B decays. Each of the tagging and mistag parameters is varied in turn by the uncertainty from the correction. The deviations obtained are added in quadrature to give the quoted uncertainty from this source.
- Allowing for possible CP violation in the transverse polarization, and in the WT longitudinally polarized signal SCF events results in additional uncertainties on signal observables. We vary S and C by ± 0.5 (± 1.0) for the transverse polarization (WT SCF).
- Possible CP violation from interference in doubly Cabibbo-suppressed decays (DCSD) on the tag side of the event [53] contribute to systematic uncertainties on S_{long} and C_{long} .
- We estimate the systematic error on our results coming from neglecting the interference between $B^0 \rightarrow \rho^+ \rho^-$ and other 4π final states: $B \rightarrow a_1 \pi$, $\rho \pi \pi^0$ and $\pi \pi \pi^0 \pi^0$. Details of this calculation are given in the Appendix.

- As the PDFs used in the ML fit do not account for all of the correlations between discriminating variables used in the fit, the results have a small bias. We calculate the fit bias on the signal observables as described in Section VIII. The observed bias on N_{signal} , and f_L is corrected, and 100% of the correction is assigned as a systematic uncertainty. We do not observe a significant bias on S_{long} and C_{long} , and conservatively assign a systematic uncertainty from fit bias based on the statistical precision of this test.
- Small imperfections in the knowledge of the geometry of the SVT over time can affect the measurement of S_{long} and C_{long} . We vary the alignment according to the results obtained from the study of $e^+e^- \rightarrow e^+e^-$, $\mu^+\mu^-$ events in order to estimate the magnitude of this systematic error on our CP results.

The branching fraction has multiplicative systematic uncertainties from the reconstruction of π^0 mesons in the detector (6%), uncertainties in the reconstruction of charged particles (0.8%), and the discrimination of $\pi^\pm$ from other types of charged particles (1%). In addition to these uncertainties, there is a 1.1% uncertainty on the number of $B\bar{B}$ pairs in the data sample. The statistical uncertainties arising from the MC samples used in this analysis are negligible.

X. CONSTRAINTS ON THE UNITARITY TRIANGLE

A. The $SU(2)$ isospin analysis

In $SU(2)$ isospin symmetry, the amplitudes of B decays to $\rho\rho$ final states [20] are related by:

$$\frac{1}{\sqrt{2}}A^{+-} = A^{+0} - A^{00}, \quad (28)$$

$$\frac{1}{\sqrt{2}}\bar{A}^{+-} = \bar{A}^{-0} - \bar{A}^{00}, \quad (29)$$

for the longitudinal polarization and each of the CP eigenstates of the transverse polarization, where A^{ij} ($\bar{A}^{ij}$) are the amplitudes of $B(\bar{B})$ decays to the final state with charge ij , where $i = +, -, 0$ and $j = -, 0$. These two relations correspond to triangles in a complex plane as shown in Fig. 8. In the usual phase convention [20], the amplitudes $\bar{A}$ are rotated to $\bar{A}$ in order to align the base of the triangles. After aligning A^{+0} and $\bar{A}^{-0}$, the phase difference between A^{+-} and $\bar{A}^{+-}$ is $2\delta\alpha$. There are two sources of ambiguities on the measurement of α . There is a four-fold ambiguity coming from the orientation of the isospin triangles, and a two-fold ambiguity from the trigonometric conversion in Eq. (12). In order to measure α one must measure the branching fractions and charge asymmetries of B decays to $\rho^+\rho^-$, $\rho^\pm\rho^0$, $\rho^0\rho^0$.




FIG. 8: The isospin triangle for $B \rightarrow \rho\rho$ decays.

There are several assumptions implicitly used in the isospin-based direct measurements of α :

- This approach only considers tree and gluonic penguin contributions. Possible contributions from electroweak penguins (EWP) are neglected as they do not obey $SU(2)$ isospin symmetry. These EWPs have the same topology as the gluonic penguin diagram in Fig. 2, with the gluon replaced by γ or Z^0 bosons. In the absence of EWP contributions $|A^{+0}| = |\tilde{A}^{-0}|$, and A_{CP} is zero for $B^+ \rightarrow \rho^+\rho^0$. Several groups have estimated the correction due to the $SU(2)$ breaking effect of EWP contributions to be $1.5 - 2.0$ degrees [54, 55]. These estimates consider contributions from the two EWP operators assumed to be dominant in the effective Hamiltonian.
- The possible effect on the isospin analysis from $\rho^0 - \omega$ mixing [55] is neglected.
- Other $SU(2)$ symmetry breaking effects are neglected. Estimates of the magnitude of these effects are much less than the current experimental precision [56, 57]. Possible isospin 5/2 amplitudes also break the $SU(2)$ triangle construction [58].
- The isospin analysis outlined above neglects possible $I=1$ amplitudes [59]. The presence of $I=1$ amplitudes in $B^0 \rightarrow \rho^+\rho^-$ can be tested by measuring S_{long} and C_{long} for different ranges of the invariant $\pi^\pm\pi^0$ mass.

We constrain the CKM angle α and penguin contribution $\delta\alpha$ from an isospin analysis of $B \rightarrow \rho\rho$ decays. The inputs to the isospin analysis are the amplitudes of the CP -even longitudinal polarization of the $\rho\rho$ final state, as well as the measured values of S_{long} and C_{long} for $B^0 \rightarrow \rho^+\rho^-$. We use the following numerical inputs in the isospin analysis:

- The measurements of $\mathcal{B}(B^0 \rightarrow \rho^+\rho^-)$, f_L , S_{long} , and C_{long} presented here.
- The measurements of $\mathcal{B}$ and f_L for $B^+ \rightarrow \rho^+\rho^0$ from Ref. [21].
- The measurement of $\mathcal{B}(B^0 \rightarrow \rho^0\rho^0)$ from Ref. [23].




FIG. 9: Confidence level on α obtained from the isospin analysis with the statistical method described in [14]. The dashed lines correspond to the 68% (top) and 90% (bottom) CL intervals.

To interpret our results in terms of a constraint on α from the isospin relations, we construct a χ^2 that includes the measured quantities expressed as the lengths of the sides of the isospin triangles and we determine the minimum χ_0^2 . We have adopted a simulated-experiment technique to compute the confidence level (CL) on α ; our method is similar to the approach proposed in Ref. [60]. For each value of α , scanned between 0 and 180° , we determine the difference $\Delta\chi_{\text{DATA}}^2(\alpha)$ between the minimum of $\chi^2(\alpha)$ and χ_0^2 . We then generate MC experiments around the central values obtained from the fit to data with the given value of α and we apply the same procedure. The fraction of these experiments in which $\Delta\chi_{\text{MC}}^2(\alpha)$ is smaller than $\Delta\chi_{\text{DATA}}^2(\alpha)$ is interpreted as the CL on α . Figure 9 shows $1 - \text{CL}$ for α obtained from this method.

It is possible to obtain a prediction of α from indirect constraints by combining measurements of the CKM matrix elements $|V_{us}|$, $|V_{ud}|$, $|V_{ub}|$, and $|V_{cb}|$, CP violation in mixing from neutral kaons, B - $\bar{B}$ mixing in B_d and B_s mesons and the measurement of $\sin 2\beta$ from $b \rightarrow c\bar{c}s$ decays. The indirect constraint on α from the UTfit [15] and CKMfitter [14] groups are $(92.9 \pm 5.7)^\circ$ and $(100^{+4.5}_{-7.3})^\circ$ respectively. Recent calculations using QCD Factorization predict $\alpha = (85.6^{+7.4}_{-7.3})^\circ$ [61]. Selecting the solution closest to the CKM combined fit average we find $\alpha = [73.1, 117.0]^\circ$ at 68% CL, where the error is dominated by $|\delta\alpha|$ which is 18° at 68% CL. The constraint obtained on α is worse than the previous result in Ref. [26] for the following reasons: (i) the central value of the branching fraction for $B^+ \rightarrow \rho^+\rho^0$ [21] is smaller than the value previously used [22, 62], and (ii) the results of the latest search for $B^0 \rightarrow \rho^0\rho^0$ show evidence for a signal [23], while the previous constraint on α used the central value corresponding to the upper limit available at that time [63]. Both of these factors lead to an increase in the penguin contribution to the total uncertainty on

α when using an isospin analysis.

A future extension to the $SU(2)$ isospin method to measure α described here will be possible when there are sufficient statistics to perform a time-dependent analysis of $B^0 \rightarrow \rho^0 \rho^0$. The parameters S_{long} and C_{long} measured in $B^0 \rightarrow \rho^0 \rho^0$ can be incorporated into the isospin analysis, allowing one to over-constrain the isospin triangles of Eq. (28) and Eq. (29).

B. Flavor $SU(3)$ analysis

There has been progress on constraining the penguin contribution in $B^0 \rightarrow \rho^+ \rho^-$ using a flavor $SU(3)$

based approach and experimental constraints from $B^+ \rightarrow K^{*0} \rho^+$ [24]. The amplitude for $B^0 \rightarrow \rho^+ \rho^-$ can be written as

$$\mathcal{A}(B^0 \rightarrow \rho^+ \rho^-) = T e^{i\gamma} + P e^{i\delta_{\text{TP}}}, \quad (30)$$

where T and P are the magnitudes of tree and penguin amplitudes, γ is the phase of V_{ub} , and δ_{TP} is the strong phase difference between T and P . The approach of Ref. [24] relates the penguin amplitude in $B^0 \rightarrow \rho^+ \rho^-$ to the amplitude of the penguin decay $B^+ \rightarrow K^{*0} \rho^+$ giving three relations:

$$C_{\text{long}} = \frac{2r \sin \delta_{\text{TP}} \sin(\beta + \alpha)}{1 - 2r \cos \delta_{\text{TP}} \cos(\beta + \alpha) + r^2}, \quad (31)$$

$$S_{\text{long}} = \frac{\sin 2\alpha + 2r \cos \delta_{\text{TP}} \sin(\beta - \alpha) - r^2 \sin 2\beta}{1 - 2r \cos \delta_{\text{TP}} \cos(\beta + \alpha) + r^2}, \quad (32)$$

$$\left(\frac{|V_{cd}|f_\rho}{|V_{cs}|f_{K^*}} \right)^2 \frac{\Gamma_L(B^\pm \rightarrow K^{*0} \rho^\pm)}{\Gamma_L(B^0 \rightarrow \rho^+ \rho^-)} = \frac{F r^2}{1 - 2r \cos \delta_{\text{TP}} \cos(\beta + \alpha) + r^2}, \quad (33)$$

with three unknowns, α , r , and δ_{TP} . The parameter $r = |P/T|$, and β is the phase of V_{td} . We use the value of β obtained from $b \rightarrow c \bar{c} s$ decays [7, 8]. The CP averaged decay rates of the longitudinal components of $B^\pm \rightarrow K^{*0} \rho^\pm$ and $B^0 \rightarrow \rho^+ \rho^-$ (Γ_L) are related by the squared ratio of CKM matrix elements and decay constants (f_i , where ρ or K^*) as shown in Eq. (33). The factor F accounts for additional sources of $SU(3)$ breaking not described by the decay constants.

The assumptions used in this approach are:

- The amplitude in the penguin dominated decay $B^+ \rightarrow K^{*0} \rho^+$ is related to the penguin amplitude in $B^0 \rightarrow \rho^+ \rho^-$,
- The dominant $SU(3)$ breaking correction accounted for by F is the neglect of annihilation diagrams in the $B^+ \rightarrow K^{*0} \rho^+$ decay. We use the same value of F as Ref. [24]; $F = 0.9 \pm 0.6$.

The result is a constraint on α with theoretical uncertainty from penguin contributions of $(+2, -5)^\circ$ which represents a stronger constraint on the unitarity triangle than provided by the isospin analysis constraint. Figure 10 shows $1 - \text{CL}$ for α obtained with this method. Selecting the solution closest to the CKM combined fit average [14, 15] we find $\alpha = [83.3, 105.8]^\circ$ at 68% CL, where the error is dominated by the experimental uncertainty on S_{long} and C_{long} . The strong phase difference δ_{TP} is only weakly constrained, and $r = 0.10_{-0.04}^{+0.03}$. The QCD

factorization approach in Refs. [64–66] predicts that δ_{TP} is small, and so we require $|\delta_{\text{TP}}| < 90^\circ$ as in Ref. [24]. We find $\alpha = (89.8_{-6.4}^{+7.0})^\circ$. These constraints are in agreement with the prediction for α from UTfit [15] and CKMfitter [14], these predictions are $(92.9 \pm 5.7)^\circ$ and $(100_{-7.3}^{+4.5})^\circ$, respectively.




FIG. 10: Confidence level on α obtained from the flavor $SU(3)$ approach. The horizontal dashed lines correspond to the 68% (top) and 90% (bottom) CL intervals. The constraint shown in this figure uses some of the same inputs as the $SU(2)$ isospin constraint of Fig. 9, and the two can not be averaged.

XI. CONCLUSIONS

We report the measurement of the branching fraction, f_L , and CP violation parameters, S_{long} and C_{long} , for the decay $B^0 \rightarrow \rho^+ \rho^-$ using a data sample of $(383.6 \pm 4.2) \times 10^6 B\bar{B}$ pairs. We obtain the following results:

$$\begin{aligned} \mathcal{B}(B^0 \rightarrow \rho^+ \rho^-) &= (25.5 \pm 2.1(\text{stat})^{+3.6}_{-3.9}(\text{syst})) \times 10^{-6}, \\ f_L &= 0.992 \pm 0.024(\text{stat})^{+0.026}_{-0.013}(\text{syst}), \\ S_{\text{long}} &= -0.17 \pm 0.20(\text{stat})^{+0.05}_{-0.06}(\text{syst}), \\ C_{\text{long}} &= 0.01 \pm 0.15(\text{stat}) \pm 0.06(\text{syst}). \end{aligned}$$

Using these results, and experimental knowledge of the other $B \rightarrow \rho\rho$ final states, we perform an isospin analysis to obtain a measurement of the CKM angle α . The confidence level distribution for α is shown in Fig. 9. The solution obtained that is compatible with the results of SM-based fits of existing data is $\alpha = [73.1, 117.0]^\circ$ at 68% CL. The uncertainty on the measurement of α from the isospin analysis is dominated by penguin pollution. If one uses the flavor $SU(3)$ approach described in the text to constrain α , one obtains the constraint $\alpha = [83.3, 105.8]^\circ$ at 68% CL.

XII. ACKNOWLEDGMENTS

We are grateful for the extraordinary contributions of our PEP-II colleagues in achieving the excellent luminosity and machine conditions that have made this work possible. The success of this project also relies critically on the expertise and dedication of the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and the kind hospitality extended to them. This work is supported by the US Department of Energy and National Science Foundation, the Natural Sciences and Engineering Research Council (Canada), the Commissariat à l’Energie Atomique and Institut National de Physique Nucléaire et de Physique des Particules (France), the Bundesministerium für Bildung und Forschung and Deutsche Forschungsgemeinschaft (Germany), the Istituto Nazionale di Fisica Nucleare (Italy), the Foundation for Fundamental Research on Matter (The Netherlands), the Research Council of Norway, the Ministry of Science and Technology of the Russian Federation, Ministerio de Educación y Ciencia (Spain), and the Science and Technology Facilities Council (United Kingdom). Individuals have received support from the Marie-Curie IEF program (European Union) and the A. P. Sloan Foundation.

APPENDIX: INTERFERENCE OF $B^0 \rightarrow \rho^+ \rho^-$ WITH OTHER $B^0 \rightarrow 4\pi$ MODES

An extensive study of the B backgrounds, and the associated systematic error, has been presented earlier in

this paper. A number of B backgrounds decay into the same final state as the signal. These are:

- $B^0 \rightarrow a_1^\pm \pi^\mp, a_1^\pm \rightarrow \rho^\pm \pi^0$,
- $B^0 \rightarrow \rho^\mp \pi^\pm \pi^0$, non resonant,
- $B^0 \rightarrow \pi^+ \pi^- \pi^0 \pi^0$, non resonant.

The systematic error associated with these modes was evaluated by propagating the uncertainty on the branching ratio and CP asymmetries to the final measurements, with the appropriate description of acceptance, SCF, resolutions and other reconstruction biases (see Section IX B). The likelihood described in Section VII does not account for possible interference between $B^0 \rightarrow \rho^+ \rho^-$ and other $B^0 \rightarrow 4\pi$ final states.

Acceptance, SCF, and other reconstruction biases are not taken into account in this study, and instead, the $\rho^+ \rho^-$ measurements are averaged in a perfect region of the phase space referred to as the “ $\rho^+ \rho^-$ band”, on which the analysis technique is assumed to have a uniform sensitivity. This “ $\rho^+ \rho^-$ band” corresponds to the kinematic selection of $\rho^\pm$ (see Section III B): $0.5 < m_{\pi^\pm \pi^0} < 1.0 \text{ GeV}/c^2$ and $-0.9 < \cos \theta_i < 0.98$.

The remainder of this Appendix describes the final state wave function, decay amplitudes, effective CP asymmetries and the estimate of the systematic uncertainty from neglecting interference between the signal and other $B^0 \rightarrow 4\pi$ final states.

1. The final state wave functions

This section summarizes the kinematic dependence of the decay amplitudes.

a. Lorentz invariant phase space

The four-particle final state can be described completely in terms of the following five variables:

- m_1 and m_2 : the masses of the $\pi^+ \pi^0$ and $\pi^- \pi^0$ pairs
- $\cos \theta_1$ and $\cos \theta_2$: the angle between the $\pi^\pm \pi^0$ pair line of flight (as seen in the B frame) and the π^0 in the pion pair frame.
- ϕ : the angle between the two planes defined by the $\pi^+ \pi^0$ and $\pi^- \pi^0$ pairs in the B frame.

We write each phase space dependent quantity $d\Phi_4$ as a function of these five variables. These are:

$$d\Phi_4 \propto \frac{P_B^{\pi^+ \pi^0}}{M_B} \frac{P_{\pi^+ \pi^0}^{\pi^0}}{m_1} \frac{P_{\pi^- \pi^0}^{\pi^0}}{m_2} dm_1^2 dm_2^2 d\cos \theta_1 d\cos \theta_2 d\phi, \quad (34)$$

where $P_B^{\pi^+ \pi^0}$ denotes the momentum of the pion pair in the B frame and M_B is the mass of the B (and similarly for the other quantities in this expression).

b. Kinematic dependence of $\rho^+\rho^-$

We assume that the longitudinal polarization of $\rho^+\rho^-$ is dominant, so the kinematic dependence of the $\rho^+\rho^-$ wave function is:

$$f(\rho^+\rho^-) \propto BW(m_1)BW(m_2)Y_1^0(\theta_1)Y_1^0(\theta_2) \propto BW(m_1)BW(m_2)\cos(\theta_1)\cos(\theta_2), \quad (35)$$

where BW denotes a Breit-Wigner function (the simple non-relativistic form is assumed), and Y_l^m are the spherical harmonics.

c. Kinematic dependence of $a_1\pi$

In order to write the wave function of $a_1^+\pi^-$, we need to define the following three variables, which are functions of the variables found in part 1a of this Appendix:

- θ_ρ : the angle between the π^0 and the ρ line of flight (as seen from the a_1) in the ρ frame,
- θ_{a_1} : the angle between the ρ and a_1 line of flight (as seen from the B) in the a_1 frame,
- ϕ_{a_1} : the angle between the two decay planes of the a_1 and ρ mesons, in the a_1 frame.

A schematic view of $B^0 \rightarrow a_1^\pm \pi^\mp$ showing θ_ρ and θ_{a_1} is given in Fig. 11.




FIG. 11: Definition of the variables θ_ρ and θ_{a_1} entering the $a_1\pi$ kinematic dependence.

The a_1 has the possibility to decay both into an S wave and a D wave. We assume that the S wave contribution is dominant, so the kinematic dependence of $a_1\pi$ is:

$$f(a_1\pi) \propto BW(a_1)BW(\rho) \left\{ d_{0,1}^1(\theta_{a_1})Y_1^1(\theta_\rho, \phi_{a_1}) + d_{0,0}^1(\theta_{a_1})Y_1^0(\theta_\rho) + d_{0,-1}^1(\theta_{a_1})Y_1^{-1}(\theta_\rho, \phi_{a_1}) \right\}, \quad (36)$$

where the $d_{m,m'}^j$ are rotation matrices. A non-relativistic Breit-Wigner is assumed for $BW(\rho)$, and a relativistic form is used for $BW(a_1)$.

d. Kinematic dependence of the non-resonant contributions

The kinematic dependence of the non-resonant contributions can be written as:

$$f(\rho^\mp \pi^\pm \pi^0) \propto BW(\rho_1)Y_1^0(\theta_1), \quad (37)$$

for $\rho^\mp \pi^\pm \pi^0$, and:

$$f(\pi^+ \pi^- \pi^0 \pi^0) = \text{constant}, \quad (38)$$

for the non-resonant 4-pion final state.

We note that the spherical harmonics entering the $\rho^+\rho^-$ amplitudes found in Eq. (35) differ from those found in these two amplitudes: this means that these wave functions are orthogonal, and after the integration over the helicity angles, the interfering terms $f(\rho^+\rho^-)f^*(\rho^\mp \pi^\pm \pi^0)$ and $f(\rho^+\rho^-)f^*(\pi^+ \pi^- \pi^0 \pi^0)$ will cancel completely if the integration is done over the whole phase space. The “ $\rho^+ \rho^-$ band” used in this analysis is slightly asymmetric in helicity angle, and as a result we can expect a small deviation from zero for these integrals.

e. Symmetrization of the wave functions

The final state is made of four bosons, two of which are identical (the two π^0). Thus each of the above wave functions needs to be symmetrized, that is to say that the amplitude:

$$f(p_{\pi^+}, p_{\pi^0}^1, p_{\pi^-}, p_{\pi^0}^2), \quad (39)$$

where p denotes the four-momentum of one of the pions, needs to be replaced by:

$$\tilde{f}(p_{\pi^+}, p_{\pi^0}^1, p_{\pi^-}, p_{\pi^0}^2) = \frac{1}{\sqrt{2}} \left\{ f(p_{\pi^+}, p_{\pi^0}^1, p_{\pi^-}, p_{\pi^0}^2) + f(p_{\pi^+}, p_{\pi^0}^2, p_{\pi^-}, p_{\pi^0}^1) \right\}. \quad (40)$$

In the rest of this study, each of the wave functions are symmetrized by π^0 meson exchange.

2. Decay amplitudes

For a particular contribution $B^0 \rightarrow x$, the total decay amplitude $\mathcal{A}(B^0 \rightarrow x)$ is the product of a phase space independent amplitude $A(B^0 \rightarrow x)$ and one of the above kinematic dependences $\tilde{f}(x)$:

$$\mathcal{A}(B^0 \rightarrow x) = A(B^0 \rightarrow x)\tilde{f}(x). \quad (41)$$

This section describes how we calculate $A(B^0 \rightarrow x)$.

a. Amplitude of $B \rightarrow \rho^+ \rho^-$

For $B \rightarrow \rho^+ \rho^-$, two amplitudes need to be known: $A(B^0 \rightarrow \rho^+ \rho^-)$ and $A(\bar{B}^0 \rightarrow \rho^+ \rho^-)$, corresponding to 4 degrees of freedom. We choose the following parameterization:

- The branching ratio $\mathcal{B}(B \rightarrow \rho^+ \rho^-)$,
- The CP -violating asymmetries $C_{\text{long}}(\rho^+ \rho^-)$ and $S_{\text{long}}(\rho^+ \rho^-)$,
- An overall phase $\delta(\rho^+ \rho^-)$ which may be set to zero in the appropriate phase convention.

The parameters $\mathcal{B}(B \rightarrow \rho^+ \rho^-)$, $C_{\text{long}}(\rho^+ \rho^-)$ and $S_{\text{long}}(\rho^+ \rho^-)$ are set to the values reported here (see Section XI).

b. Amplitude of $B \rightarrow a_1 \pi$

Four amplitudes describe the $B \rightarrow a_1 \pi$ contributions, corresponding to a total of 8 degrees of freedom: $A(B^0 \rightarrow a_1^+ \pi^-)$, $A(B^0 \rightarrow a_1^- \pi^+)$, $A(\bar{B}^0 \rightarrow a_1^+ \pi^-)$ and $A(\bar{B}^0 \rightarrow a_1^- \pi^+)$. We choose a parameterization similar to that of the time dependent $a_1 \pi$ analysis [67], for which parameters have direct physical meaning:

- $\mathcal{B}(B^0 \rightarrow a_1 \pi)$ is the total branching ratio,
- $C(a_1 \pi) + \Delta C(a_1 \pi) = C^+(a_1 \pi) = \frac{1 - |\lambda_{CP}(a_1^+ \pi^-)|^2}{1 + |\lambda_{CP}(a_1^+ \pi^-)|^2}$
where $\lambda_{CP}(a_1^+ \pi^-) = \frac{q}{p} \frac{A(\bar{B}^0 \rightarrow a_1^+ \pi^-)}{A(B^0 \rightarrow a_1^+ \pi^-)}$,
- $C(a_1 \pi) - \Delta C(a_1 \pi) = C^-(a_1 \pi) = \frac{1 - |\lambda_{CP}(a_1^- \pi^+)|^2}{1 + |\lambda_{CP}(a_1^- \pi^+)|^2}$
where $\lambda_{CP}(a_1^- \pi^+) = \frac{q}{p} \frac{A(\bar{B}^0 \rightarrow a_1^- \pi^+)}{A(B^0 \rightarrow a_1^- \pi^+)}$, $C(a_1 \pi)$ is a direct CP -violating asymmetry and $\Delta C(a_1 \pi)$ is CP -conserving,
- $S(a_1 \pi) + \Delta S(a_1 \pi) = S^+(a_1 \pi) = \frac{2\text{Im}\{\lambda_{CP}(a_1^+ \pi^-)\}}{1 + |\lambda_{CP}(a_1^+ \pi^-)|^2}$,
- $S(a_1 \pi) - \Delta S(a_1 \pi) = S^-(a_1 \pi) = \frac{2\text{Im}\{\lambda_{CP}(a_1^- \pi^+)\}}{1 + |\lambda_{CP}(a_1^- \pi^+)|^2}$,
 $S(a_1 \pi)$ is a CP -violating asymmetry in the interference between decay and mixing, and $\Delta S(a_1 \pi)$ is CP conserving,
- $A_{CP}(a_1 \pi) = \frac{|A^{+-}|^2 + |\bar{A}^{+-}|^2 - |A^{-+}|^2 - |\bar{A}^{-+}|^2}{|A^{+-}|^2 + |\bar{A}^{+-}|^2 + |A^{-+}|^2 + |\bar{A}^{-+}|^2}$ where $A^{\pm\mp} = A(B^0 \rightarrow a_1^\pm \pi^\mp)$, and $\bar{A}^{\pm\mp}$ is its complex

conjugate. These are also direct CP violating asymmetries,

- Two overall phases of the final states $\delta(a_1^+ \pi^-)$ and $\delta(a_1^- \pi^+)$.

For $a_1 \pi$ we assume that all of these parameters are constant over the whole of phase space.

3. Effective CP asymmetries

Considering the two final states $\rho^+ \rho^-$ and $a_1 \pi$, the phase-space dependent four-pion decay amplitude is:

$$\begin{aligned} \mathcal{A}(B^0 \rightarrow 4\pi) &= \mathcal{A}(B^0 \rightarrow \rho^+ \rho^-) + \mathcal{A}(B^0 \rightarrow a_1^+ \pi^-) \\ &\quad + \mathcal{A}(B^0 \rightarrow a_1^- \pi^+), \\ &= A(B^0 \rightarrow \rho^+ \rho^-) f(\rho^+ \rho^-) \\ &\quad + A(B^0 \rightarrow a_1^+ \pi^-) f(a_1^+ \pi^-) \\ &\quad + A(B^0 \rightarrow a_1^- \pi^+) f(a_1^- \pi^+), \end{aligned}$$

with a similar expression for $\mathcal{A}(\bar{B}^0 \rightarrow 4\pi)$. Similar equations exist for $\rho\pi\pi$ and 4π final states.

We can define the phase-space dependent quantity:

$$\lambda_{4\pi}(\Phi_4) = \frac{q}{p} \frac{\mathcal{A}(\bar{B}^0 \rightarrow 4\pi)}{\mathcal{A}(B^0 \rightarrow 4\pi)}, \quad (42)$$

where $\lambda_{4\pi}(\Phi_4)$ is a function of m_1 , m_2 , $\cos\theta_1$, $\cos\theta_2$, and ϕ . Thus define the phase-space dependent CP -violating asymmetries:

$$C_{4\pi}(\Phi_4) = \frac{1 - |\lambda_{4\pi}(\Phi_4)|^2}{1 + |\lambda_{4\pi}(\Phi_4)|^2}, \quad (43)$$

$$S_{4\pi}(\Phi_4) = \frac{2\text{Im}(\lambda_{4\pi}(\Phi_4))}{1 + |\lambda_{4\pi}(\Phi_4)|^2}, \quad (44)$$

that are analogous to Eq. (8) and Eq. (9).

In principle it is possible to measure the value of $C_{4\pi}(\Phi_4)$ and $S_{4\pi}(\Phi_4)$ at each point of phase space, however this would be experimentally challenging. Instead, this measurement uses integrated values C_{eff} and S_{eff} , which can be seen as the averages of C and S across the previously defined “ $\rho^+ \rho^-$ band”, weighted over the number of events found at each location of the phase space which corresponds to:

$$\begin{aligned}
C_{\text{eff}} &= \frac{\int_{\rho\rho \text{ band}} C_{4\pi}(\Phi_4)(|\mathcal{A}(B^0 \rightarrow 4\pi)|^2 + |\mathcal{A}(\bar{B}^0 \rightarrow 4\pi)|^2)d\Phi_4}{\int_{\rho\rho \text{ band}} (|\mathcal{A}(B^0 \rightarrow 4\pi)|^2 + |\mathcal{A}(\bar{B}^0 \rightarrow 4\pi)|^2)d\Phi_4}, \\
S_{\text{eff}} &= \frac{\int_{\rho\rho \text{ band}} S_{4\pi}(\Phi_4)(|\mathcal{A}(B^0 \rightarrow 4\pi)|^2 + |\mathcal{A}(\bar{B}^0 \rightarrow 4\pi)|^2)d\Phi_4}{\int_{\rho\rho \text{ band}} (|\mathcal{A}(B^0 \rightarrow 4\pi)|^2 + |\mathcal{A}(\bar{B}^0 \rightarrow 4\pi)|^2)d\Phi_4}.
\end{aligned} \tag{45}$$

We can also define the values of C and S that would be observed in the absence of interference effects, denoted by $C_{4\pi}^{\mathcal{I}}(\Phi_4)$ and $S_{4\pi}^{\mathcal{I}}(\Phi_4)$, and similarly the integrated values across the “ $\rho^+\rho^-$ band” $C_{\text{eff}}^{\mathcal{I}}$ and $S_{\text{eff}}^{\mathcal{I}}$. Finally, for the evaluation of the systematic error, we are really interested in the biases on C and S purely due to interference which are the two differences $\delta C = C_{\text{eff}} - C_{\text{eff}}^{\mathcal{I}}$ and $\delta S = S_{\text{eff}} - S_{\text{eff}}^{\mathcal{I}}$. We adopt the same strategy for the branching fraction $\mathcal{B}(\rho^+\rho^-)$ by defining $\delta\mathcal{B} = \mathcal{B}_{\text{eff}} - \mathcal{B}_{\text{eff}}^{\mathcal{I}}$ in a similar way.

4. The systematic error

As the 8 parameters describing the amplitude $A(B^0(\bar{B}^0) \rightarrow a_1\pi)$ are not yet measured, to estimate the systematic error on $\rho^+\rho^-$ parameters, we need to propagate the uncertainty on these 8 parameters. For this, we use a MC technique. For each experiment, we generate the CP parameters $C(a_1\pi)$, $\Delta C(a_1\pi)$, $S(a_1\pi)$, $\Delta S(a_1\pi)$, $A_{CP}(a_1\pi)$, the phases $\delta(a_1^+\pi^-)$ and $\delta(a_1^-\pi^+)$, and the branching fraction $\mathcal{B}(a_1\pi)$. The values used to generate the $a_1\pi$ contribution have Gaussian PDFs for measured quantities (with mean and width corresponding to the measurement), and uniform PDFs for unmeasured quantities (sampled over the allowed values). Then, we estimate the value of δC and δS for each generated value of the $a_1\pi$ parameters. The systematic error are taken as the RMS of the δC , δS and $\delta\mathcal{B}$ distributions.

For this study, the quantity $\mathcal{B}(a_1\pi)$ is fixed to the value measured in [51], $\mathcal{B}(B^0 \rightarrow a_1^+\pi^-, a_1^+ \rightarrow \rho^+\pi^0) = (16.6 \pm 2.4) \times 10^{-6}$ and the CP parameters are obtained from [67]: $C(a_1\pi) = -0.10 \pm 0.17$, $\Delta C(a_1\pi) = 0.26 \pm 0.17$, $S(a_1\pi) = 0.37 \pm 0.22$, $\Delta S(a_1\pi) = -0.14 \pm 0.22$ and $A_{CP}(a_1\pi) = -0.07 \pm 0.07$.

We estimate the effect of interference between signal and either $\rho^\mp\pi^\pm\pi^0$ or $\pi^+\pi^-\pi^0\pi^0$ in a similar way, where we assume the branching fraction for these two non-resonant backgrounds is $(5.7 \pm 13.0) \times 10^{-6}$. This value is obtained by modeling the $\rho^\mp\pi^\pm\pi^0$ component in the likelihood fit and by fitting a free yield to the data, and we assume that $\rho^\mp\pi^\pm\pi^0$ and $\pi^+\pi^-\pi^0\pi^0$ have the same branching fraction.

We summarize the systematic uncertainty obtained for the different 4π final states considered here in Table VI. The total uncertainty on $\mathcal{B}$ is used to calculate the quoted systematic uncertainty on f_L from this source. We con-

clude that interference between signal and other 4π final states has a negligible impact on the measurement.

TABLE VI: Systematic error coming from interference effects due to three contaminating amplitudes.

Contribution	$\sigma(C)$	$\sigma(S)$	$\frac{\sigma(\mathcal{B})}{\mathcal{B}}$
$a_1\pi$	0.009	0.009	2.5%
$\rho\pi\pi$ non resonant	0.010	0.010	1.1%
4π non resonant	0.002	0.002	0.3%
Total	0.014	0.014	2.7%

-
- [1] J. H. Christenson, J. W. Cronin, V. L. Fitch, and R. Turlay. *Phys. Rev. Lett.* **13**, 138 (1964).
- [2] A. D. Sakharov. *Pisma Zh. Eksp. Teor. Fiz.*, **5**, 32 (1967); Traduction JETP Lett **5**, 24 (1967).
- [3] N. Cabibbo. *Phys. Rev. Lett.* **10**, 531 (1963).
- [4] M. Kobayashi and T. Maskawa. *Prog. Theor. Phys.* **49**, 652 (1973).
- [5] L. Wolfenstein. *Phys. Rev. Lett.* **51**, 1945 (1983).
- [6] A. J. Buras, M. E. Lautenbacher, and G. Ostermaier. *Phys. Rev. D* **50**, 3433 (1994).
- [7] B. Aubert *et al.* (BABAR Collaboration). *Phys. Rev. Lett.* **89**, 201802 (2002).
- [8] K. Abe *et al.* (Belle Collaboration). *Phys. Rev. D* **66**, 071102 (2002).
- [9] M. B. Gavela. *Mod. Phys. Lett. A* **9**, 795 (2002).
- [10] P. Huet and E. Sather. *Phys. Rev. D* **51**, 379 (1995).
- [11] B. Aubert *et al.* (BABAR Collaboration). hep-ex/0703021.
- [12] K.-F. Chen *et al.* (Belle Collaboration). *Phys. Rev. Lett.* **98**, 031802 (2007).
- [13] M. Ciuchini, E. Franco, G. Martinelli, L. Reina, and L. Silvestrini. *Z. Phys. C* **68**, 239 (1995).
- [14] CKMfitter Group. *Eur. Phys. J. C* **41**, 1-131 (2005). Updated results and plots available at: <http://ckmfitter.in2p3.fr>.
- [15] UTfit Collaboration. *JHEP* **0507**, 028 (2005). Updated results and plots available at: <http://www.utfit.org>.
- [16] I. Bigi and A. Sanda. *CP Violation*. Cambridge University Press, 1999.
- [17] G. Branco, L. Lavoura, and J. Silva. *CP Violation*. Oxford University Press, 1999.
- [18] Unless explicitly stated, charge conjugation is implied throughout.
- [19] A. Bevan. *Mod. Phys. Lett. A* **21**, 305 (2006).
- [20] M. Gronau and D. London. *Phys. Rev. Lett.* **65**, 3381 (1990).
- [21] B. Aubert *et al.* (BABAR Collaboration). *Phys. Rev. Lett.* **97**, 261801 (2006).
- [22] J. Zhang *et al.* (Belle Collaboration). *Phys. Rev. Lett.* **91**, 221801 (2003).
- [23] B. Aubert *et al.* (BABAR Collaboration). *Phys. Rev. Lett.* **98**, 111801 (2007).
- [24] M. Beneke, M. Gronau, J. Rohrer, and M. Spranger. *Phys. Lett. B* **638**, 68 (2006).
- [25] See, for example, J. Zupan hep-ph/0701004, and references therein.
- [26] B. Aubert *et al.* (BABAR Collaboration). *Phys. Rev. Lett.* **95**, 041805 (2005).
- [27] A. Somov *et al.* (Belle Collaboration). *Phys. Rev. Lett.* **96**, 171801 (2006).
- [28] W.-M. Yao *et al.* (Particle Data Group). *J. Phys.* **G33**, 1 (2006).
- [29] See, for example, D. Kirkby and Y. Nir, Section 12, in Ref. [28].
- [30] B. Aubert *et al.* (BABAR Collaboration). *Phys. Rev. Lett.* **93**, 231801 (2004).
- [31] A. Somov *et al.* (Belle Collaboration). hep-ex/0702009.
- [32] PEP-II Conceptual Design Report. *SLAC-R-418*, (1993).
- [33] B. Aubert *et al.* (BABAR Collaboration). *Nucl. Instrum. Methods Phys. Res., Sect. A* **479**, 1 (2002).
- [34] G. Benelli, K. Honscheid, E. A. Lewis, J. J. Regensburger, and D. S. Smith. *Nuclear Science Symposium Conference Record, 2005 IEEE*, **2**, 1145 (2005).
- [35] W. Menges. *Nuclear Science Symposium Conference Record, 2005 IEEE*, **3**, 1470 (2005).
- [36] M. R. Convery, P. C. Kim, H. P. Paar, C. H. Rogers, R. H. Schindler, S. K. Swain, and C. C. Young. *Nucl. Instrum. Methods Phys. Res., Sect. A* **556**, 134 (2006).
- [37] S. Agostinelli *et al.* (GEANT4 Collaboration). *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [38] A. Drescher, B. Grawe, B. Hahn, B. Ingelbach, U. Matthiesen, H. Scheck, J. Spengler, and D. Wegener. *Nucl. Instrum. Methods Phys. Res., Sect. A* **237**, 464 (1985).
- [39] G. C. Fox and S. Wolfram. *Phys. Lett.* **41**, 1581 (1978).
- [40] B. Aubert *et al.* (BABAR Collaboration). *Phys. Rev. D* **66**, 032003 (2002).
- [41] M. J. Oreglia. Ph.D Thesis, SLAC-236, Appendix D, (1980).
- [42] J. E. Gaiser. Ph.D Thesis, SLAC-255, Appendix F, (1982).
- [43] T. Skwarnicki. Ph.D Thesis, DESY F31-86-02, Appendix E, (1986).
- [44] H. Albrecht *et al.* (ARGUS Collaboration). *Phys Lett*, B241:278, (1990).
- [45] B. Aubert *et al.* (BABAR Collaboration). *Phys. Rev. D* **72**, 072003 (2005).
- [46] A. Garmash *et al.* (Belle Collaboration). *Phys. Rev. Lett.* **96**, 251803 (2006).
- [47] B. Aubert *et al.* (BABAR Collaboration). hep-ex/0701035.
- [48] J. Zhang *et al.* (Belle Collaboration). *Phys. Rev. Lett.* **94**, 031801 (2005).
- [49] CLEO Collaboration. *Phys. Rev. Lett.* **85**, 2881 (2000).
- [50] B. Aubert *et al.* (BABAR Collaboration). *Phys. Rev. Lett.* **91**, 201802 (2003).
- [51] B. Aubert *et al.* (BABAR Collaboration). *Phys. Rev. Lett.* **97**, 051802 (2006).
- [52] B. Aubert *et al.* (BABAR Collaboration). *Phys. Rev. D* **74**, 031104 (2006).
- [53] O. Long, M. Baak, R. N. Cahn, and D. Kirkby. *Phys. Rev. D* **68**, 034010 (2003).
- [54] See, for example, p73 of Ref. [14].
- [55] M. Gronau and J. Zupan. *Phys. Rev. D* **71**, 074017 (2005).
- [56] S. Gardner. hep-ph/9906269.
- [57] S. Gardner. *Phys. Rev. D* **72**, 034015 (2005).
- [58] F. J. Botella, D. London, and J. P. Silva. *Phys. Rev. D* **73**, 071501 (2006).
- [59] A. F. Falk, Z. Ligeti, Y. Nir, and H. Quinn. *Phys. Rev. D* **69**, 011502 (2004).
- [60] G. Feldman and R. Cousins. *Phys. Rev. D* **57**, 3873 (1998).
- [61] M. Beneke, J. Rohrer, and D. Yang. hep-ph/0612290.
- [62] B. Aubert *et al.* (BABAR Collaboration). *Phys. Rev. Lett.* **91**, 171802 (2003).
- [63] B. Aubert *et al.* (BABAR Collaboration). *Phys. Rev. Lett.* **94**, 131801 (2005).
- [64] M. Beneke, G. Buchalla, M. Neubert, and C. T. Sachrajda. *Phys. Rev. Lett.* **83**, 1914 (1999).
- [65] M. Beneke, G. Buchalla, M. Neubert, and C. T. Sachrajda. *Nucl. Phys. B* **591**, 313 (2000).

- [66] M. Beneke, G. Buchalla, M. Neubert, and C. T. Sachrajda. *Nucl. Phys. B* **606**, 245 (2001).
- [67] B. Aubert *et al.* (BABAR Collaboration). *Phys. Rev. Lett.* **98**, 181803 (2007).